

Quark-Hadron Duality and Scaling in Reduced QCD

Zoltán Batiz¹ and Franz Gross^{2,3}

¹*Centro de Física das Interações Fundamentais (CFIF),
Instituto Superior Técnico, P-1049-001 Lisboa, Portugal*

²*Department of Physics, College of William and Mary, Williamsburg, Virginia 23185*

³*Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606*

(Dated: February 8, 2008)

Abstract

We introduce a generalization of $1+1$ dimensional large N_c QCD, which we refer to as “reduced” QCD, or rQCD. In this model gluons and quark momenta live in $1+1$ dimensions only, but the quark spin and all other particles (leptons, and the photon) live in the full $1+3$ dimensions. The bound states of quarks and antiquarks are identical to those originally described by ’t Hooft (except that there are new transversely polarized states previously excluded), so the model is exactly solvable. However, significant differences arise when the model is applied to electromagnetic interactions. After reviewing the strongly interacting sector of the theory, we discuss deep inelastic scattering (DIS) in this model, and show that the new states with transverse polarization give the Callan-Gross relation and remove the pathological features of the original $1+1$ dimensional description. We conclude that rQCD gives a satisfactory description of the phenomenology and provides a deep understanding of both duality and DIS.

PACS numbers: 12.39.-x, 11.10.St, 13.40.Gp

I. INTRODUCTION

In 1974, ’t Hooft described the behavior of the large N_c limit of QCD in $1+1$ dimensions [1]. He showed that the model confined quarks and was exactly solvable. His work was followed quickly by several papers [2, 3, 4] discussing various aspects of the model. The work by Einhorn [3] discussed the ’t Hooft model predictions for deep inelastic scattering (DIS). Einhorn found that the DIS cross section was proportional to the square of the undressed quark masses, and hence would approach zero as the quark masses approach zero. This pathological result made it difficult to apply the ’t Hooft model to the phenomenology of DIS; clearly something major was missing. Of course, limiting the physics to $1+1$ dimensions also excludes transverse spin degrees of freedom, so W_1 was zero and the Callan-Gross relation [5] did not hold.

Since then QCD in $1+1$ dimensions has been studied from a variety of points of view [6, 7, 8, 9, 10] and applied recently to the study of duality in heavy meson decays [11, 12, 13]. However, to the best of our knowledge, no one has followed up on Einhorn’s study [3] of DIS.

A few years ago we found that a very simple toy model of scalar and spinor fields in $1+1$ dimension gave nice phenomenological results for DIS [14]. By extending the spinor degrees of freedom to $1+3$ dimensions we were able to recover the Callan-Gross relation. We decided to extend our approach to the 't Hooft model, and learned of the pathology in its description of DIS. It turns out that this pathology can be removed by extending the spinor degrees of freedom to $1+3$ dimensions, just as we did before, and this paper grew out of that study.

This paper is divided into five sections, with seven appendices that include many of the details. Following this brief introduction, Sec. II reviews the 't Hooft model results for the strong sector. Our discussion focuses on (i) a method of treatment of the confining interaction that removes all singularities from the theory and gives finite dressed quark masses, and (ii) the consequence of including transverse degrees of freedom for the spin of the quarks. We calculate the dressed quark mass, and the properties and spectrum of $q\bar{q}$ bound states, and also study the consequences of the completeness of the bound states. We show how to construct an off-shell $q\bar{q}$ scattering matrix, and show that the on-shell scattering matrix must be zero, as required for confined particles. Sections III and IV study electromagnetic interactions and deep inelastic scattering. Here the presence of transverse spin degrees of freedom completely alters the Einhorn discussion, giving us phenomenologically useful results. We conclude with a brief discussion.

In a subsequent paper [15] we plan to present a numerical study of the approach to scaling in the DIS limit.

II. THE 'T HOOFT MODEL

A. The Lagrangian and confining interaction

We propose that the action have the following form

$$\mathcal{S} = \mathcal{S}_{QCD} + \mathcal{S}_{QED} \quad (2.1)$$

where the QCD part of the action (gluons and quark momenta or coordinates) live in $1+1$ dimensions only, but the quark spin and the QED part (leptons and the photon) live in the full $1+3$ dimensions

$$\begin{aligned} \mathcal{S}_{QCD} &= \int dt dz \mathcal{L}_{QCD}(t, z) \\ \mathcal{S}_{QED} &= \int dt d\mathbf{r} \mathcal{L}_{QED}(t, \mathbf{r}). \end{aligned} \quad (2.2)$$

In his original paper, 't Hooft [1] discussed the strong interactions only, and in this section we review the $1+1$ dimensional model of QCD that he presented. The QED part of the action (2.2), and reduced QCD (rQCD), will be discussed in Sec. III.

The QCD Lagrangian density is

$$\mathcal{L}_{QCD}(t, z) = -\frac{1}{4} \text{Tr} [F^{\mu\nu} F_{\mu\nu}] + \sum_i \bar{q}_i (iD_\mu \gamma^\mu - m_{0i}) q_i, \quad (2.3)$$

where $q_i = q_i(t, z)$ is the quark field with flavor i and bare mass m_{0i} . The gluon field quantities are

$$A^\mu = \frac{1}{2} A_a^\mu \lambda_a \quad (2.4)$$

$$\begin{aligned}
F_{\mu\nu} &= \partial_\mu A_\nu - \partial_\nu A_\mu + ig_0 [A_\mu, A_\nu] \\
D_\mu &= \partial_\mu + ig_0 A_\mu,
\end{aligned}$$

where $F_{\mu\nu}$ is the gluon field tensor and $A_a^\mu = A_a^\mu(t, z)$ are the gluon fields with the Lorentz index μ and the color index a . The gluon fields have components in the 0 and 3 direction only, so the sum over the index μ is restricted to 0 and 3. The QCD coupling constant is g_0 and λ^a are the generators of the $SU(N_c)$ color group, normalized to

$$\begin{aligned}
\text{trace} [\lambda^a \lambda^b] &= 2 \delta^{ab} \\
\sum_a \lambda^a \lambda^a &= \frac{2(N_c^2 - 1)}{N_c} \mathbb{1},
\end{aligned} \tag{2.5}$$

where matrix multiplication of the λ matrices is implied. As $N_c \rightarrow \infty$, finite results are obtained if $g_0 \rightarrow 0$ as

$$g_0 = \frac{g}{\sqrt{2N_c}}. \tag{2.6}$$

where the effective coupling g is a constant.

While the space-time coordinates, momenta, and Lorentz vector sums in Eq. (2.3) are restricted to 1+1 dimension, we assume that the gamma matrices have the usual 4×4 Dirac structure. The quark fields are therefore a direct product of four Dirac dimensions and N_c color dimensions.

Note that the action (2.2) is Lorentz invariant under the subgroup G_3 of Lorentz transformations that leave the xy plane invariant. Specifically, the group is generated by the hamiltonian $\mathcal{H}$, and the boost $\mathcal{K}_3$, momentum $\mathcal{P}_3$, and angular momentum $\mathcal{J}_3$, operators that generate boosts and translations along the z axis, and rotations about the z axis. Four of the six commutation relations between these generators are zero, and the other two close

$$[\mathcal{K}_3, \mathcal{P}_3] = -i\mathcal{H} \quad [\mathcal{K}_3, \mathcal{H}] = -i\mathcal{P}_3, \tag{2.7}$$

guaranteeing that G_3 is a group.

Following 't Hooft [1] we introduce light cone variables:

$$b_+ = \frac{1}{\sqrt{2}} (b^0 + b^3) \tag{2.8}$$

$$b_- = \frac{1}{\sqrt{2}} (b^0 - b^3)$$

$$\mathbf{b}_\perp = \{b_1, b_2\} \tag{2.9}$$

for any arbitrary vector b (in the QCD sector, the only perpendicular components come from matrix elements of $\gamma_\perp$). Note that the scalar product of any two vectors a and b is

$$a_\mu b^\mu = a \cdot b = a_+ b_- + a_- b_+ - \mathbf{a}_\perp \cdot \mathbf{b}_\perp \tag{2.10}$$

The derivatives are defined

$$\begin{aligned}
\partial_- &= \frac{\partial}{\partial x_+} = \frac{1}{\sqrt{2}} (\partial^0 - \partial^3) = \frac{1}{\sqrt{2}} \left(\frac{\partial}{\partial x^0} + \frac{\partial}{\partial x^3} \right), \\
\partial_+ &= \frac{\partial}{\partial x_-} = \frac{1}{\sqrt{2}} (\partial^0 + \partial^3) = \frac{1}{\sqrt{2}} \left(\frac{\partial}{\partial x^0} - \frac{\partial}{\partial x^3} \right)
\end{aligned} \tag{2.11}$$

so that the divergence of a two-vector is

$$\partial_0 b^0 + \partial_3 b^3 = \partial_- b_+ + \partial_+ b_- . \quad (2.12)$$

In the same way, we can define the $+$, $-$, and $\perp$ components of the γ matrices. The anticommutation relations are all zero except for

$$\begin{aligned} \{\gamma_+, \gamma_-\} &= 2 \\ \{\gamma_x, \gamma_x\} &= \{\gamma_y, \gamma_y\} = -2 . \end{aligned} \quad (2.13)$$

Since the gluon fields are confined to 1+1 dimensions, there is only one nonvanishing component of the gluon field strength tensor

$$F_{+-} = -F_{-+} = \partial_+ A_- - \partial_- A_+ + ig_0[A_-, A_+] . \quad (2.14)$$

The QCD part of the theory is simplified if we choose the light cone gauge, where $A_- = 0$, so that the commutator contained in the field tensor F_{+-} disappears and there is only one nonzero component (A_+) of the gluon field. The Lagrangian density (2.3) then reduces to

$$\mathcal{L}_{QCD} = \frac{1}{2} Tr \left[(\partial_- A_+)^2 \right] + \sum_i \bar{q}_i (i\partial_+ \gamma_- + i\partial_- \gamma_+ - g_0 \gamma_- A_+ - m_{0i}) q_i . \quad (2.15)$$

The equation of motion for this field is then

$$\partial_-^2 A_+^a = \left(\frac{\partial}{\partial x_+} \right)^2 A_+^a = -g_0 \sum_i \bar{q}_i \lambda^a \gamma_- q_i . \quad (2.16)$$

The solution of (2.16) is

$$A_+^a(x_+, x_-) = g_0 \int dy_+ \mathcal{G}(x_+ - y_+) \sum_i \bar{q}_i(y_+, x_-) \lambda^a \gamma_- q_i(y_+, x_-) , \quad (2.17)$$

where the Green's function $\mathcal{G}$ is

$$\mathcal{G}(x_+ - y_+) = -\frac{1}{2} |x_+ - y_+| + c_1(x_+ - y_+) + c_2 . \quad (2.18)$$

The coefficients c_1 and c_2 cannot be determined without knowing the boundary conditions, so they are free parameters. The gauge condition did not eliminate all superfluous degrees of freedom, just as the Coulomb gauge, or the Lorentz gauge, do not determine uniquely the photon propagator in QED (Gribov ambiguity). We can therefore set the coefficients c_1 and c_2 equal to zero (a specific choice of gauge) in order to simplify our calculations.

Einhorn discussed the gauge issues related to these two parameters and showed that the eigenvalues of the two body bound state equation are independent of the choice of c_1 and c_2 . However, the dressed quark mass does depend on the choice of c_2 , and this in turn implies that the location of the quark mass pole is gauge dependent. Since any physically meaningful quantity is gauge invariant, we can conclude that the location of the mass pole is not physically meaningful, and this can happen only if the quarks are confined, so that free quark states do not exist. This is an indirect consequence of confinement. Direct consequences of confinement will be discussed in the next subsection when the two-body bound state equation is discussed.

Equation (2.17) shows that the gluonic field is no longer a dynamical variable, and that there are therefore no ghosts.

The Fourier transform of the Green's function (2.18) gives us the gluon “propagator”, or more precisely the momentum dependence of the effective quark-quark interaction. With c_1 and c_2 equal to zero, it can be written

$$\tilde{\mathcal{G}}(k_-) = \int_{-\infty}^{\infty} dx_+ e^{i(k_- x_+)} \mathcal{G}(x_+) \rightarrow \left\{ \frac{1}{k_-^2} - \delta(k_-) \int_{-\infty}^{\infty} \frac{d\ell_-}{\ell_-^2} \right\}. \quad (2.19)$$

This is a singular operator that is well defined only when it is part of an integral over k_- . The second term in Eq. (2.19) was introduced by Gross and Milana [16] in a different context. Its purpose is to preserve the condition

$$\mathcal{G}(0) = 0 = \int_{-\infty}^{\infty} dk_- \tilde{\mathcal{G}}(k_-) \quad (2.20)$$

and it also insures that the potential is finite for any finite value of x_+ .

The only variables remaining are the quark fields. The Feynman rule for the (undressed) quark propagator is

$$-iS_0(k) = \frac{-i}{m_0 - k_- \gamma_+ - k_+ \gamma_- - i\epsilon} = -i \frac{m_0 + k_- \gamma_+ + k_+ \gamma_-}{m_0^2 - 2k_+ k_- - i\epsilon} \quad (2.21)$$

and the quark-“gluon” coupling is

$$-i\mathcal{V}_{qqg}^a = -ig_0 \lambda^a \gamma_- , \quad (2.22)$$

leading to the following result for the exchange of gluons with momentum k_- between quarks with Dirac indices 1 and 2

$$\begin{aligned} -i\mathcal{V}_{qq} &= \sum_a \left(-i\mathcal{V}_{qqg}^{a1} \right) \mathcal{O}_{12} \left(-i\mathcal{V}_{qqg}^{a2} \right) \left[-i\tilde{\mathcal{G}}(k_-) \right] \\ &= -ig_0^2 \frac{2(N_c^2 - 1)}{N_c} \gamma_-^1 \mathcal{O}_{12} \gamma_-^2 \tilde{\mathcal{G}}(k_-) \rightarrow -ig^2 \gamma_-^1 \mathcal{O}_{12} \gamma_-^2 \tilde{\mathcal{G}}(k_-), \end{aligned} \quad (2.23)$$

where we used Eq. (2.5) (with matrix multiplication of the λ matrices implied), and took the limit as $N_c \rightarrow \infty$ using the definition (2.6).

The quark self-energy and the dressed quark mass are calculated in the next subsection.

B. Dyson-Schwinger Equation for the dressed quark

We determine the dressed single quark propagator, $S(p)$, using the (one body) Dyson Schwinger equation (DSE):

$$\begin{aligned} S(p) &= S_0(p) - S(p) \Sigma(p) S_0(p) \\ &= S_0(p) - S(p) \left[-ig^2 \int \frac{d^2 k}{(2\pi)^2} \tilde{\mathcal{G}}(p_- - k_-) \gamma_- S(k) \gamma_- \right] S_0(p), \end{aligned} \quad (2.24)$$

shown graphically in Fig. 1. The gluon interaction does not mix quark flavors, and the calculation is identical for each flavor of quark, so the flavor index is suppressed. The




FIG. 1: Dyson-Schwinger equation for the quark. The corkscrew line is the gluon interaction, the thin line the undressed quark propagator, and the heavy solid line the dressed quark propagator.

rainbow approximation (undressed vertices and the absence of the quark loops from the gluon propagator) is justified in the large N_c limit [1]. Since for every internal loop there is a factor of $\alpha^2 = g^2/(2N_c)$, and a multiplicative factor of $\sum_a \lambda^a \lambda^a = 2N_c$, the color dependence disappears. The vertex corrections and the quark-gluon vertices do not have a multiplicative factor of N_c , and are therefore suppressed in the large N_c limit.

In Eq. (2.24), $d^2k = dk_- dk_+$, and since D does not depend on k_+ , it follows immediately that the self-energy integral does not depend on p_+ either, and must have the form $\Sigma(p) = B(p_-)\gamma_-$. Hence the dressed propagator is of the form

$$S(p) = \frac{1}{m_0 - p_- \gamma_+ - [p_+ - B(p_-)] \gamma_- - i\epsilon}, \quad (2.25)$$

where, using $\gamma_- \gamma_+ \gamma_- = 2\gamma_-$, the self-energy contribution is then

$$B(p_-) = -2i g^2 \int \frac{dk_- dk_+}{(2\pi)^2} \frac{k_- \tilde{\mathcal{G}}(p_- - k_-)}{m_0^2 - 2k_- (k_+ - B(k_-)) - i\epsilon}. \quad (2.26)$$

Performing the k_+ integral gives

$$\int dk_+ \frac{k_-}{m_0^2 - 2k_- (k_+ - B(k_-)) - i\epsilon} = \frac{i\pi}{2} \text{sig}(k_-), \quad (2.27)$$

and substituting this back into Eq. (2.26) gives

$$B(p_-) = \frac{g^2}{4\pi} \int dk_- \tilde{\mathcal{G}}(p_- - k_-) \text{sig}(k_-). \quad (2.28)$$

Using (2.19) for $\tilde{\mathcal{G}}(p_- - k_-)$ gives

$$\begin{aligned} B(p_-) &= \frac{g^2}{4\pi} \int_{-\infty}^{\infty} dk_- \left\{ \frac{\text{sig}(k_-)}{(p_- - k_-)^2} - \frac{\text{sig}(p_-)}{(p_- - k_-)^2} \right\} \\ &= -\frac{g^2}{2\pi p_-}. \end{aligned} \quad (2.29)$$

Substituting this back into Eq. (2.23) gives

$$S(p) = \frac{1}{m_0 - p_- \gamma_+ - \left[p_+ + \frac{g^2}{2\pi p_-} \right] \gamma_- - i\epsilon} = \frac{m_0 + p_- \gamma_+ + \left(p_+ + \frac{g^2}{2\pi p_-} \right) \gamma_-}{m_0^2 - \frac{g^2}{\pi} - 2p_+ p_- - i\epsilon}. \quad (2.30)$$

Note that the quark mass pole has been shifted to a *smaller* value

$$m_0^2 \rightarrow m^2 = m_0^2 - \frac{g^2}{\pi}, \quad (2.31)$$

where m is the dressed quark mass. It turns out that choosing $c_1 \neq 0$ will not effect the dressing of the mass, but choosing $c_2 \neq 0$ would give

$$m^2 = m_0^2 - \frac{g^2}{\pi} (1 - \pi c_2 |p_-|). \quad (2.32)$$

This gives a mass that is momentum dependent and not covariant. Hence the dressed mass is gauge dependent and unphysical. With these cautionary remarks, we choose $c_1 = c_2 = 0$ because the mass is covariant. One undesirable feature of this choice is that $m^2 < 0$ in the chiral limit. This could be corrected by choosing c_2 large and positive.

Having obtained the dressed propagator, we are able to proceed with the two body bound state calculation.

C. Two-body bound states

1. Algebraic form of the two body equations

Consider a bound state of a $q\bar{q}$ pair. The quark has dressed mass m_1 and electric charge e_1 , and the antiquark (which might be of a different flavor) has dressed mass m_2 and charge e_2 . The momentum of the bound state is r , the momentum of the quark is p and the momentum of the outgoing antiquark is $r - p$, as shown in Fig. 2. In much of the following discussion, we will treat the outgoing antiquark as an incoming quark with momentum $p - r$, and following this convention label the bound state vertex function $\Gamma(p, p - r)$. For a color neutral state we may carry out the color sums using (2.23), giving the Bethe-Salpeter equation for the bound state vertex function:

$$\Gamma(p, p - r) = i \int \frac{d^2 k}{(2\pi)^2} V(p, k) \gamma_- S_1(k) \Gamma(k, k - r) S_2(k - r) \gamma_-, \quad (2.33)$$

where S_1 and S_2 are dressed quark propagators, and the kernel is the singular operator

$$V(p, k) = g^2 \tilde{\mathcal{G}}(p_- - k_-). \quad (2.34)$$

With the substitution $\Gamma(p, p - r) = \gamma_- G(p, p - r)$ [1], Eq. (2.33) becomes:

$$G(p, p - r) = 4i \int \frac{d^2 k}{(2\pi)^2} \frac{V(p, k) k_- (k_- - r_-) G(k, k - r)}{[m_1^2 - k^2 - i\epsilon] [m_2^2 - (k - r)^2 - i\epsilon]}. \quad (2.35)$$

The formula shows that $G(p, p - r)$ does not depend on p_+ , and hence the k_+ integration can be carried out immediately.

Integrations of the variable k_+ over two quark propagators occur frequently, and are worked out in detail in Appendix A. It is convenient to introduce the momentum fractions

$$y = \frac{k_-}{r_-} \quad z = \frac{p_-}{r_-}. \quad (2.36)$$




FIG. 2: The Bethe-Salpeter equation

If the fraction z lies in the interval $[0,1]$, Eq. (2.35) reduces to

$$\begin{aligned} \mu_n^2 \Phi_n(z, r) &= \left(\frac{\alpha_1 + 1}{z} + \frac{\alpha_2 + 1}{1 - z} \right) \Phi_n(z, r) - \oint_0^1 dy \frac{\Phi_n(y, r) - \Phi_n(z, r)}{(z - y)^2} \\ &\equiv \frac{\pi}{g^2} H(z) \Phi_n(z, r) \quad (\text{if } z \in [0, 1]) , \end{aligned} \quad (2.37)$$

where $\oint$ is the principal value integral, and μ_n , α_1 , and α_2 are dimensionless parameters

$$\mu_n^2 = \frac{\pi M_n^2}{g^2}, \quad \alpha_1 = \frac{\pi m_1^2}{g^2}, \quad \alpha_2 = \frac{\pi m_2^2}{g^2}, \quad (2.38)$$

with $r^2 = M_n^2$ the anticipated mass eigenvalues that will emerge from the solution of the equation, and m_i the *dressed* quark masses. The two-body “wave function” $\Phi_n(z, r)$, is defined by

$$\Phi_n(z, r) = \frac{\mathcal{N} r_- G_n(z, r)}{\Delta(z, M_n^2)} = \frac{\mathcal{N} r_- G_n(z, r)}{\left(\frac{m_1^2}{z} + \frac{m_2^2}{1 - z} - M_n^2 \right)} = \frac{\mathcal{N} r_- G_n(z, r)}{\frac{g^2}{\pi} \left(\frac{\alpha_1}{z} + \frac{\alpha_2}{1 - z} - \mu_n^2 \right)}, \quad (2.39)$$

where $G_n(z, r) = G(p, p - r)$ is the anticipated eigenfunction solution for the vertex corresponding to the n th bound state, and $\mathcal{N}$ is a normalization constant. It is chosen so that

$$1 = \int_0^1 dz \Phi_n^2(z, r), \quad (2.40)$$

and will be calculated in subsection III C. Note that Eq. (2.37) defines the two-quark Hamiltonian, $H(z)$, on the interval $0 \leq z \leq 1$. The *bare* quark masses enter into the Hamiltonian, while the *dressed* quark masses enter into the relation (2.39) between the vertex function and the wave function.

If the momentum fraction z lies *outside* of the interval $0 \leq z \leq 1$, the vertex function defined by Eq. (2.35) is not zero, but can be obtained by quadrature from the vertex function defined *inside* of the interval $[0,1]$. Using (A8) and the definition (2.39) gives

$$\Phi_n(z, r) = \frac{1}{\left(\frac{\alpha_1}{z} + \frac{\alpha_2}{1 - z} - \mu_n^2 \right)} \int_0^1 dy \frac{\Phi_n(y, r)}{(z - y)^2} \quad (\text{if } z \notin [0, 1]) . \quad (2.41)$$

These definitions and equations will be used in the following sections when the hadronic currents are computed.

An alternative form of Eq. (2.37) follows if we integrate the last term by parts:

$$\begin{aligned} \mu_n^2 \Phi_n(z, r) = & \left(\frac{\alpha_1 + 1}{z} + \frac{\alpha_2 + 1}{1 - z} \right) \Phi_n(z, r) + \int_0^1 dy \frac{\Phi'_n(y, r)}{(z - y)} \\ & + \left(\frac{\Phi_n(0, r) - \Phi_n(z, r)}{z} \right) + \left(\frac{\Phi_n(1, r) - \Phi_n(z, r)}{1 - z} \right), \end{aligned} \quad (2.42)$$

where $\Phi'_n(y, r) = \partial \Phi(y, r) / \partial y$. If $\Phi_n(0, r) = \Phi_n(1, r) = 0$, which is true in all but the chiral limit, this form of the equation follows directly from 't Hooft's principle value prescription [1], so that our two-body equation is identical with 't Hooft's, even though we have finite dressed quark masses.

To avoid confusion later, we call attention to the fact that the wave function defined by Eqs. (2.39) and (2.41) is not identical to the Bethe-Salpeter wave function. The latter can be defined

$$\Psi(p, r) \equiv \frac{\mathcal{N} p_-(p - r)_- G(p, p - r)}{(m_1^2 - p^2 - i\epsilon)(m_2^2 - (p - r)^2 - i\epsilon)}, \quad (2.43)$$

where the factor $p_-(p - r)_-$ in the numerator comes from the γ_- factors in the quark propagators S . This wave function carries the p_+ dependence of the quark propagators. If we integrate (2.43) over p_+ (using the fact that G does not depend on p_+) we obtain

$$\psi(p_-, r) \equiv \int_{-\infty}^{\infty} dp_+ \Psi(p, r) = \begin{cases} \frac{\mathcal{N} r_- G(p, p - r)}{\Delta(z, r^2)} = \Phi(z, r) & \text{if } z \in [0, 1] \\ 0 & \text{if } z \notin [0, 1] \end{cases}. \quad (2.44)$$

This is the “equal t_- ” wave function, and equals the wave function we are using *only in the region* $z \in [0, 1]$. As we have seen, the wave function $\Phi(z, r)$ that solves the 't Hooft equation is non zero *for all* z .

We conclude this discussion of the two body wave equation by returning to Eq. (2.33). In place of the replacement $\Gamma(p, p - r) = \gamma_- G_n(z, r)$, it could equally well have been assumed that

$$\Gamma(p, p - r) = \gamma_- \gamma_k G_n(z, r), \quad (2.45)$$

where k is either x or y (recall that we are assuming that the Dirac matrices live in 1+3 dimensional space). Note that the Bethe-Salpeter equations with the substitutions (2.45) reduce to the same equation (2.35) for G_n , and hence *each two-body state is triply degenerate, with spin structure given by* γ_- , $\gamma_- \gamma_x$, or $\gamma_- \gamma_y$. In each of these three cases the wave functions and bound state mass are identical. The additional states with the γ_x or γ_y structure are outside the scope of the original 't Hooft model, but must be considered in reduced QCD. They are states with quark spins in the transverse direction, and will play a major role in the discussion of DIS below.

The chiral limit is exactly soluable and of some interest. If $m_{0i} = 0$, then Eq. (2.37) has the normalized ground state solution

$$M_0 = 0, \quad \Phi_0(z, r) = 1. \quad (2.46)$$

Outside of the interval $[0, 1]$, Eq. (2.41) gives the same result, so that

$$\Phi_0(z, r) = 1 \quad (2.47)$$

for all z in $(-\infty, \infty)$.




FIG. 3: Diagrammatic representation of confinement via two-body interaction. The $\times$ indicates that the dressed quark is on-shell, and at the point when both quarks are on-shell, the vertex function is zero.

2. Confinement in the 't Hooft model

The 't Hooft model illustrates how confinement can be realized in two (apparently different) ways. If the singularities of the confining interaction are left unregularized, as they were in 't Hooft's original paper [1], the quark masses are infinite. Even if the singularity is regularized as in this paper, whenever $m_{0i}^2 < g^2/\pi$ the dressed quark mass will be unphysical, and there is no real quark mass pole. This insures that the quarks are confined.

In the second case ($m_{0i}^2 > g^2/\pi$ and $m_i^2 > 0$) the dressed quark mass is real and positive and there is a quark mass pole. What keeps the quarks confined in this case? The answer is that the quark vertex function $G_n(z, r)$ has a *zero* at precisely the value of z where all of the quarks in the bound state could be on mass shell. This requirement is illustrated diagrammatically in Fig. 3, and seems miraculous. However, as discussed in Ref. [17], it is only a consequence of fact that the *wave function* is finite at the pole (because the wave equation permits no singularities), and therefore the vertex function

$$G_n(z, r) = \left[\frac{m_1^2}{z} + \frac{m_2^2}{1-z} - M_n^2 \right] \frac{\Phi(z, r)}{\mathcal{N}r_-} \quad (2.48)$$

must be zero. Precisely the same mechanism works in the Schrödinger description of two nonrelativistic particles bound by a linear (or any other) confining potential.

In summary, quark mass poles are always unphysical because the dressed mass is gauge dependent, and bound state poles are physical because they are gauge invariant. In this respect the undressed quark mass is the only physically significant parameter. The two ways of viewing confinement are therefore not exclusive, but equivalent, since they are related through gauge transformations.

3. Numerical solutions for meson wave functions

The bound state equation was solved numerically using modified cubic splines. The splines and the technique are fully described in Appendix G. Each spline is a smooth function (continuous up to its third derivative) and has support over a small region of the z axis. The splines overlap, so that by independently adjusting the height of each spline it is possible to smoothly approximate any function defined on the interval $[0, 1]$.

We have modified the standard spline technique to insure that the numerical solutions satisfy the boundary conditions exactly. These boundary conditions are determined by examining Eq. (2.37) in the vicinity of $z \sim 0$ and $z \sim 1$ [1, 11], and require the wave function

TABLE I: Numerical values of M_n^2 for $m_{01} = m_{02} = 1.5$ and various n_s (number of splines) and n_g (number of guass points for each spline segment).

level	n_s	20			40			60		
n	n_g	2	4	6	2	4	6	2	4	6
1		13.730	13.732	13.732	13.717	13.718	13.718	13.714	13.715	13.715
2		25.613	25.614	25.615	25.572	25.573	25.573	25.563	25.564	25.564
10		108.95	109.35	109.36	108.42	108.43	108.43	108.28	108.28	108.28
20		186.17	210.60	210.58	208.82	209.69	209.69	208.88	208.97	208.97
30		–	–	–	299.64	314.02	314.02	308.04	309.38	309.38
40		–	–	–	362.04	409.77	409.74	410.31	411.24	411.24

TABLE II: Numerical values of M_n^2 for $m_{01} = m_{02} = 0.5$ and various n_s and n_g .

level	n_s	20			40			60		
n	n_g	4	6	8	4	6	8	4	6	8
1		2.6196	2.6625	2.6841	2.6485	2.6786	2.6936	2.6616	2.6858	2.6979
2		10.540	10.650	10.706	10.616	10.692	10.731	10.648	10.710	10.741
10		86.170	86.371	86.477	86.055	86.288	86.409	86.197	86.400	86.503
20		199.44	199.43	199.43	183.93	184.11	184.21	183.53	183.77	183.90
30		–	–	–	287.16	287.22	287.25	282.09	282.27	282.36
40		–	–	–	399.18	399.15	399.15	383.13	383.22	383.27

go like

$$\begin{aligned}
\Phi_n(z, r) &\rightarrow z^{\beta_1} & (\text{as } z \rightarrow 0) & \text{ with } \pi\beta_1 = -\alpha_1 \tan(\pi\beta_1) \\
\Phi_n(z, r) &\rightarrow (1-z)^{\beta_2} & (\text{as } z \rightarrow 1) & \text{ with } \pi\beta_2 = -\alpha_2 \tan(\pi\beta_2).
\end{aligned} \tag{2.49}$$

Note that the $\beta_i = 0$ in the chiral limit (when $\alpha_i = -1$), consistent with the exact solution (2.47). To fit these boundary conditions the splines closest to $z = 0$ and $z = 1$ have these fractional powers built into their functional form, as described in Appendix G.

The wave function is expanded into n_s splines, each with four segments (except for those at the boundary, which have only three). The splines overlap, and taking into account that there must be at least three splines (one regular spline in the center and one modified spline at each end), the total number of segments is $n_s + 1$, dividing the interval into segments of length $1/(n_s + 1)$. The equation is turned into a matrix equation by integrating over each segment using n_g gaussian points. Tables I and II show the numerical stability. If the quark mass is larger than unity, better than 1% accuracy is achieved with only two gauss points per interval, provided $n \lesssim n_s$. As the level number n increases above $\sim n_s/2$ toward n_s , the reliability of the calculation decreases. To study highly excited states, it is sufficient to use $n_g = 2$, and push n_s as high as possible.

The situation is somewhat different if the bare quark masses are less than unity (Table II). In this case the wave function is very steep at the boundaries and 4 or 6 gaussian points per interval are needed to get 1% accuracy for the low lying states. In a subsequent paper [15], we will limit our numerical discussion of duality to cases with $m_{0i} > 1$ where the solutions are very stable, and there are no poles in the region outside of $[0,1]$ [see the brief remarks following Eq. (F14)].




FIG. 4: The square of the bound state mass (in units of g^2/π) versus the order of the state n , computed using 60 splines. The straight line is the linear relation $n \pi g^2$. Note that the masses follow this linear relation up to near $n \sim n_s = 60$, where the departure from linearity is due to numerical inaccuracy.




FIG. 5: The ground state wave functions for equal bare quark masses m_0 of 1.1, 1.5, and 3 (the wave functions become steeper at $z = 0$ and $z = 1$ as the quark mass decreases). In all cases 60 splines were used, and the wave function was normalized to a maximum value of unity.

Figure 4 shows the linearly rising Regge trajectory characteristic of the 't Hooft model. Figures 5-7 show the ground, first excited and 10th level for three cases with equal quark masses of different values. Note the “tails” of the wave functions outside of the region $[0,1]$, given by Eq. (2.41). These tails are large for small quark masses, but shrink quickly as the quark mass increases, and are also small for highly excited states. Figure 8 shows the 20th level for spline numbers 20, 40, and 60. Note that the state is not fully described even with 40 splines, supporting the observation that an accurate description of states of order n requires a spline number of approximately $\sim 2n$.




FIG. 6: The second state (first excited state) for the same three quark masses shown in Fig. 5. (See caption to Fig. 5.)




FIG. 7: The tenth state for the same three quark masses shown in Fig. 5. (See caption to Fig. 5.) In this figure the dots are $\sin(10\pi z)$, showing that this is an excellent approximation for small quark masses.

D. Scattering of quarks and antiquarks in the presence of confinement

1. Completeness relation and the two-body Greens function

The Hamiltonian defined in Eq. (2.37) is hermitian on the interval $[0, 1]$ and it is straightforward to show that the different solutions are orthogonal on this interval. With the normalization given in Eq. (2.40)

$$\int_0^1 dz \Phi_n(z, r_n) \Phi_m(z, r_m) = \delta_{nm}. \quad (2.50)$$




FIG. 8: The 20th state for and equal quark mass of 1.5. The small amplitude curve was calculated for $n_s = 20$, and completely fails to represent the state. The dotted and full amplitude solid lines are calculated for $n_s = 40$ and 60 , respectively. Note that $n_s = 40$ does well except for the first and last oscillation.

The completeness of the eigenfunctions implies that

$$\sum_n \Phi_n(z, r_n) \Phi_n(z', r_n) = \delta(z - z') \quad \text{if } z, z' \in [0, 1]. \quad (2.51)$$

The Greens function is therefore

$$G(z, z', q^2) = \sum_n \frac{\Phi_n(z, r_n) \Phi_n(z', r_n)}{q^2 - M_n^2}, \quad (2.52)$$

where q is any external momentum. The completeness relation shows that G has the property of a Greens function

$$\begin{aligned} \{q^2 - H(z)\} G(z, z', q^2) &= \sum_n (q^2 - H(z)) \frac{\Phi_n(z, r_n) \Phi_n(z', r_n)}{q^2 - M_n^2} \\ &= \delta(z - z') \quad \text{if } z, z' \in [0, 1]. \end{aligned} \quad (2.53)$$

The same argument also gives

$$G(z, z', q^2) \{q^2 - H(z')\} = \delta(z - z') \quad \text{if } z, z' \in [0, 1]. \quad (2.54)$$

2. The scattering matrix

In preparation for the study of electromagnetic interactions, it is useful to introduce the $q\bar{q}$ scattering matrix, defined by the following infinite series

$$\begin{aligned} M(p', p; q) \gamma_{1-} \otimes \gamma_{2-} &= V(p', p) \gamma_{1-} \otimes \gamma_{2-} \\ &+ i \int \frac{d^2 k}{(2\pi)^2} V(p', k) V(k, p) [\gamma_- S_1(k) \gamma_-]_1 \otimes [\gamma_- S_2(k - q) \gamma_-]_2 \end{aligned}$$

$$\begin{aligned}
& +i^2 \int \frac{d^2 k'}{(2\pi)^2} \int \frac{d^2 k}{(2\pi)^2} V(p', k') V(k', k) V(k, p) \\
& \quad \times [\gamma_- S_1(k') \gamma_- S_1(k) \gamma_-]_1 \otimes [\gamma_- S_2(k - q) \gamma_- S_2(k' - q) \gamma_-]_2 \\
& + \dots
\end{aligned} \tag{2.55}$$

The momentum are defined as for the two-body bound state, except that here q is the unconstrained momentum of the $q\bar{q}$ pair and q^2 is not an eigenvalue. The kernel V is the singular operator (2.34). The singularities of this operator will never be exposed because we will limit application of this scattering equation to cases where we integrate over either the initial or final momentum. Note that the structure of the equation anticipates the reduction of the dressed propagators

$$[\gamma_- S_i(k) \gamma_-]_i = \frac{k_- [\gamma_- \gamma_+ \gamma_-]_i}{m_i^2 - 2k_- k_+} = \frac{2k_-}{m_i^2 - 2k_- k_+} \gamma_{i-}, \tag{2.56}$$

so that the factor of $\gamma_{1-} \otimes \gamma_{2-}$ is common, and the series (2.55) reduces to

$$\begin{aligned}
M(p', p; q) = & V(p', p) + i \int \frac{d^2 k}{\pi^2} \frac{V(p', k) V(k, p)}{d_2(k_-, k_+, q)} \\
& + i^2 \int \frac{d^2 k'}{\pi^2} \int \frac{d^2 k}{\pi^2} \frac{V(p', k') V(k', k) V(k, p)}{d_2(k'_-, k'_+, q) d_2(k_-, k_+, q)} + \dots,
\end{aligned} \tag{2.57}$$

where d_2 was defined in Eq. (A3). From this result we can conclude that M does not depend on p'_+ and p_+ .

In applications below, we will encounter this scattering series in the following form

$$\begin{aligned}
\langle MG\mathcal{O} \rangle(p'; q) = & \int \frac{d^2 p}{\pi^2} \frac{M(p', p; q) \mathcal{O}(p_-, q)}{d_2(p_-, p_+, q)} = \int \frac{d^2 p}{\pi^2} \frac{V(p', p) \mathcal{O}(p_-, q)}{d_2(p_-, p_+, q)} \\
& + i \int \frac{d^2 k}{\pi^2} \frac{V(p', k)}{d_2(k_-, k_+, q)} \int \frac{d^2 p}{\pi^2} \frac{V(k, p) \mathcal{O}(p_-, q)}{d_2(p_-, p_+, q)} \\
& + i^2 \int \frac{d^2 k'}{\pi^2} \int \frac{d^2 k}{\pi^2} \frac{V(p', k') V(k', k)}{d_2(k'_-, k'_+, q) d_2(k_-, k_+, q)} \int \frac{d^2 p}{\pi^2} \frac{V(k, p) \mathcal{O}(p_-, q)}{d_2(p_-, p_+, q)},
\end{aligned} \tag{2.58}$$

where $\mathcal{O}(p_-, q)$ is an operator that, by assumption, does not depend on p_+ . In Appendix C we show how to use the Greens function to write this series as

$$\int_0^1 dz M(z', z; q^2) \frac{\mathcal{O}(z, q^2)}{\Delta(z, q^2)} = \sum_n \Delta(z', M_n^2) \frac{\Phi_n(z', r_n)}{M_n^2 - q^2} \int_0^1 dz \Phi_n(z, r_n) \mathcal{O}(z, q^2), \tag{2.59}$$

where we replaced $(q_-)^2 M(p', p; q)/\pi \rightarrow M(z', z; q^2)$ and $\mathcal{O}(p_-, q) \rightarrow \mathcal{O}(z, q^2)$.

This equation has a nice physical interpretation. It displays the “scattering amplitude” as the sum over the propagation of the bound states, which is all confinement will allow (without meson decay mechanisms, which are ignored in this paper, no cuts are possible). The only singularities in q^2 that can occur are poles at the bound state masses M_n^2 .

With these tools in place we are able to study the electromagnetic interaction of hadrons.

III. REDUCED QCD AND THE ELECTROMAGNETIC SECTOR

We return to the discussion of the Lagrangian density introduced in Eq. (2.2) and look at the electromagnetic sector. In rQCD the electromagnetic fields (and the electron currents) are extended to 1+3 dimensions. The Lagrangian density is therefore

$$\mathcal{L}_{QED}(t, \mathbf{r}) = -\frac{1}{4}F_0^{\mu\nu}(t, \mathbf{r})F_{0\mu\nu}(t, \mathbf{r}) + \sum_i e_i \bar{q}_i(t, z)\gamma_\mu q_i(t, z) A_0^\mu(t, \mathbf{r}) - e J_\mu^e(t, \mathbf{r}) A_0^\mu(t, \mathbf{r}), \quad (3.1)$$

where $A_0(t, \mathbf{r})$ is the electromagnetic field (with the subscript 0 distinguishing it from the gluon field), the vector sums are now over 4 dimensions with $\mu = \{0, 1, 2, 3\}$, e_i is the electromagnetic charge of quarks with flavor i , $J_\mu^e(t, \mathbf{r})$ is the electron current that produces the electromagnetic field, and

$$F_0^{\mu\nu} = \partial^\mu \mathcal{A}_0^\nu - \partial^\nu \mathcal{A}_0^\mu, \quad (3.2)$$

is the electromagnetic field tensor. Because the electron current exists in four dimensions, all four components of the electromagnetic field will be non-zero, in general, and all four components of the quark current will be excited by electromagnetic scattering.

We now apply this discussion to electron scattering, where $Q^2 = -q^2 > 0$, so that $q_- < 0$. The initial particle (quark or hadron) has momentum $p = p' - q$ and the final particle momentum p' , with momentum fractions

$$z' = \frac{p'_-}{q_-} \quad \frac{p_-}{q_-} = z' - 1. \quad (3.3)$$

If the final particle is on shell, then $p'_- > 0$ and $z' < 0$, and $z' \rightarrow -\infty$ as $Q^2 \rightarrow 0$. We will use the notation j_0^μ to denote the *bare* quark current operator, which is

$$j_0^\mu = \gamma^\mu. \quad (3.4)$$

While the dressing of the strong quark-gluon vertex is suppressed by the large N_c limit [1], the quark electromagnetic vertex can be dressed by gluon exchanges. The longitudinal part of the dressed electromagnetic vertex (j^μ) can be computed directly from the Ward-Takahashi identity

$$q_\mu j^\mu(p', p) = S^{-1}(p) - S^{-1}(p'), \quad (3.5)$$

where $p + q = p'$. Using the dressed propagator (2.30), we find that

$$\begin{aligned} j_- &= \gamma_- \\ j_+ &= \gamma_+ - \gamma_- \frac{g^2}{2\pi} \frac{1}{p'_- p_-} = \gamma_+ + \gamma_- \frac{g^2}{2\pi} \frac{1}{q_-^2 z'(1-z')}. \end{aligned} \quad (3.6)$$

Consequently the j_- component is unmodified.

A. Electromagnetic coupling to quarks and the quark form factor

It is instructive to obtain the full quark current directly from the dressing of the quark-photon vertex. The vertex is dressed by successive gluon exchanges, as shown diagrammatically in Fig. 9. When coupling to a photon, there is no flavor change, and hence, for each




FIG. 9: Diagrammatic picture of the gluonic dressing of the quark current.

flavor i , this series is

$$\begin{aligned}
 j_i^\mu(p', p' - q) &= \gamma^\mu + i \int \frac{d^2 k}{(2\pi)^2} V(p', k) \gamma_- S_i(k) \gamma^\mu S_i(k - q) \gamma_- \\
 &+ i^2 \int \frac{d^2 p}{(2\pi)^2} \int \frac{d^2 k}{(2\pi)^2} V(p', p) V(p, k) \gamma_- S_i(p) \gamma_- S_i(k) \gamma^\mu S_i(k - q) \gamma_- S_i(p - q) \gamma_- \\
 &+ \dots
 \end{aligned} \tag{3.7}$$

This series is evaluated in Appendix D using the methods developed in Appendix C. The final results are conveniently expressed in term of a quark form factor, defined to be

$$F_i(z', Q^2) \equiv \sum_n \Delta_i(z', M_n^2) \frac{\Phi_n(z', r_n)}{M_n^2 + Q^2} \int_0^1 dy \Phi_n(y, r_n), \tag{3.8}$$

where $z' = p'_-/q_-$ and $Q^2 = -q^2$ in anticipation of applications to electron scattering. Note that the form factor is expressed as a sum of bound state poles at $Q^2 = -M_n^2$, arising from the propagation of the bound states that couple to the photon. Vector dominance is a rigorous consequence of this model.

Using this form factor, the currents reduce to

$$j_{i-}(p', q) = \gamma_- + \gamma_- F_i(z', Q^2) \tag{3.9}$$

$$j_{i+}(p', q) = \gamma_+ - \gamma_- \left\{ \frac{g^2}{2\pi} \frac{1}{p'_- p_-} + \frac{q_+}{q_-} F_i(z', Q^2) \right\} \tag{3.10}$$

$$j_i^x(z', q^2) = \gamma_x - \gamma_x \gamma_- \left\{ \frac{q_-}{m_{0i}} \frac{g^2}{2\pi} \frac{1}{p'_- p_-} + \frac{q_+}{m_{0i}} F_i(z', Q^2) \right\}. \tag{3.11}$$

Note that the terms independent of the quark form factor F_i are identical to the expected result (3.6), and that the contribution from the form factor is *purely transverse*, in that

$$\left[q_+ j_{i-}(p, q) + q_- j_{i+}(p, q) \right] \Big|_{F_i \text{ term}} = \gamma_- F_i(z, Q^2) \left\{ q_+ - q_- \frac{q_+}{q_-} \right\} = 0. \tag{3.12}$$

This result mimics the method for insuring current conservation developed in Ref. [18], and gives some evidence that that method is dynamically sound.

These results are used in the next subsection to calculate the hadronic transition current.




FIG. 10: Currents for the electromagnetic transition from the initial state i to the final state f . The left hand diagram is the contribution from the electromagnetic interaction with the quark with mass m_1 and charge e_1 and the right hand diagram is the same for the antiquark with mass m_2 and charge e_2 .

B. The transition current

In the presence of confinement, and without any meson decay mechanisms, the quarks in a $q\bar{q}$ bound state must remain bound, even after they absorb an energetic photon. In this section we calculate the transition current for the process $\gamma^* + M_i \rightarrow M_f$, where M_f is the mass of a (possibly excited) final bound state and M_i is the mass of the initial (ground) state.

The transition current consists of the sum of two contributions: one in which the photon is absorbed by the quark with dressed mass m_1 , and a second in which the photon is absorbed by the antiquark with mass m_2 . The momentum used to label each of these processes are shown in Fig. 10; the initial bound state has momentum P and the final state has momentum $P_f = P + q$. The transition current coming from the first term is

$$\begin{aligned} \langle f | \mathcal{J}^\mu(P_f, P) | i \rangle \Big|_{e_1 \text{ term}} &= i e_1 N_c \int \frac{d^2 k}{(2\pi)^2} G_f(-P_f - k, -k) G_i(P + k, k) \\ &\quad \times \text{Tr} \left[\gamma_- \left\{ \begin{array}{c} 1 \\ \gamma_x \end{array} \right\} S_1(P_f + k) j_1^\mu S_1(P + k) \gamma_- S_2(k) \right] \\ &= i \int \frac{d^2 k}{(2\pi)^2} \frac{e_1 N_c G_f(-P_f - k, -k) G_i(P + k, k) N^\mu}{[m_1^2 - (P_f + k)^2 - i\epsilon][m_1^2 - (P + k)^2 - i\epsilon][m_2^2 - k^2 - i\epsilon]}, \end{aligned} \quad (3.13)$$

where j_1^μ is the quark current operator for quark 1, as worked out in subsection III A, N^μ is the spin dependent numerator discussed below, and we have allowed for the possibility that the final state may have *either* the γ_- or the $\gamma_- \gamma_x$ structure discussed above (we will not consider the $\gamma_- \gamma_y$ states; they are needed only if we consider the y component of the transverse current and give results identical to the results for the x components). Note the factor of N_c coming from the sum over all possible quark colors.

Only the term in the numerator of the S_2 propagator equal to $k_- \gamma_+$ can give a nonzero result, but the terms that contribute from the S_1 propagators depend on the matrix form of the final state, and of j_1^μ . Because the trace of a product of gamma matrices with only one factor of γ_x or γ_y is zero, the transverse currents (with γ_x or γ_y) will be zero unless the final state also includes γ_x (or γ_y for the y component of the current). Hence we have the rules:

(i) x -type transverse currents couple γ_- states to $\gamma_- \gamma_x$ states only, and (ii) longitudinal currents couple γ_- states to each other. Because of the exact degeneracy of these states, the results in both of these cases use the same momentum-space wave functions.

The integral (3.13) is evaluated in Appendix F. The result for the $-$ component of the current is

$$\begin{aligned} \langle f_- | \mathcal{J}_-(P_f, P) | i_- \rangle \Big|_{e_1 \text{ term}} &= \frac{16e_1 N_c P_-}{(2\pi) \mathcal{N}^2} \left\{ \int_y^1 + \int_0^y \mathcal{R} \right\} d\xi \Phi_f(\xi', P_f) \Phi_i(\xi, P) \\ &\quad \times \left[1 + F_1(\eta, Q^2) \right], \end{aligned} \quad (3.14)$$

where

$$y = -\frac{q_-}{P_-}, \quad \xi' = \frac{\xi - y}{1 - y}, \quad \eta = \frac{y - \xi}{y} \quad (3.15)$$

[for definitions of all the momentum fractions see Eqs. (F1) and (F2)], the state f_- has the same structure as the initial state (assumed to be γ_-), F_1 is the quark form factor defined in Eq. (3.8) and $\mathcal{R}$ is given in Eq. (F14). The plus and transverse components of the current are

$$\begin{aligned} \langle f_x | \mathcal{J}_x(P_f, P) | i_- \rangle \Big|_{e_1 \text{ term}} &= \frac{16e_1 N_c P_-}{(2\pi) \mathcal{N}^2} \left\{ \int_y^1 + \int_0^y \mathcal{R} \right\} d\xi \Phi_f(\xi', P_f) \Phi_i(\xi, P) \\ &\quad \times \frac{q_+ [m_1^2 - \eta(1 - \eta) Q^2 F_1(\eta, Q^2)]}{m_{01} \eta (1 - \eta) Q^2} \\ \langle f_- | \mathcal{J}_+(P_f, P) | i_- \rangle \Big|_{e_1 \text{ term}} &= \frac{16e_1 N_c P_-}{(2\pi) \mathcal{N}^2} \left\{ \int_y^1 + \int_0^y \mathcal{R} \right\} d\xi \Phi_f(\xi', P_f) \Phi_i(\xi, P) \\ &\quad \times \frac{q_+ [m_1^2 - \eta(1 - \eta) Q^2 F_1(\eta, Q^2)]}{q_- \eta (1 - \eta) Q^2}, \end{aligned} \quad (3.16)$$

where the state f_x has the structure $\gamma_- \gamma_x$. These results are used in the next subsection, and in the discussion of DIS.

Using the definitions of the momenta shown in Fig. 10, it is easy to see that the second term in the transition current can be obtained from the first by letting $1 \leftrightarrow 2$, as discussed in Appendix F. The exact expression for the x component of the transition current, which is the sum of the e_1 and e_2 contributions, can therefore be written

$$\langle f_x | \mathcal{J}_x(P_f, P) | i_- \rangle = \mathcal{J}_{fi}^{1,x}(Q^2, y) + \mathcal{J}_{fi}^{2,x}(Q^2, y) \quad (3.17)$$

where the reduced current $\mathcal{J}_{fi}^{j,x}(Q^2, y)$ is

$$\mathcal{J}_{fi}^{j,x}(Q^2, y) = \frac{e_j m_{0j} y}{1 - y} \left\{ \int_y^1 + \int_0^y \mathcal{R}_j \right\} d\xi \frac{\Phi_f(\xi'_j, P_f) \Phi_i(\xi_j, P)}{\xi'_j \xi_j} F_j^{\text{eff}}(\eta_j, Q^2) \quad (3.18)$$

with $\{\xi_1, \xi'_1, \eta_1\} = \{\xi, \xi', \eta\}$ and $\{\xi_2, \xi'_2, \eta_2\} = \{1 - \xi, 1 - \xi', 1 - \eta\}$ and

$$F_i^{\text{eff}}(\eta, Q^2) = \frac{1}{m_{0i}^2} \left[m_i^2 - \eta(1 - \eta) Q^2 F_i(\eta, Q^2) \right]. \quad (3.19)$$

Finally, the function $\mathcal{R}_1 = \mathcal{R}(m_1, m_2)$, was defined in Eq. (F14); the substitutions that convert the e_1 term into the e_2 term therefore give $\mathcal{R}_2 = \mathcal{R}(m_2, m_1)$.

C. Elastic scattering and normalization of the wave function

If the scattering is elastic, $x = 1$ and the momentum fraction $y = -q_-/P_-$ must be computed from the exact expressions for q and P in the c.m. frame [14]. The result, for elastic scattering from the state n , is

$$y = \frac{Q^2 + Q\sqrt{4M_n^2 + Q^2}}{2M_n^2 + Q^2 + Q\sqrt{4M_n^2 + Q^2}} \quad (3.20)$$

As $Q^2 \rightarrow 0$, $y \rightarrow 0$ and η [defined in Eq. (3.15)] approaches $-\infty$, $F_i(\eta, Q^2) \rightarrow 0$, and the *exact* result for the elastic current becomes

$$\langle n | \mathcal{J}_-(P, P) | n \rangle \Big|_{e_1 \text{ term}} = \frac{16e_1 N_c P_-}{(2\pi)\mathcal{N}^2} \int_0^1 d\xi [\Phi_n(\xi, P)]^2 = 2e_1 P_- \mathcal{F}_n(0), \quad (3.21)$$

where charge normalization requires that the form factor $\mathcal{F}_n(Q^2)$ be unity at $Q^2 = 0$. Hence

$$\mathcal{N} = 2\sqrt{\frac{N_c}{\pi}}, \quad (3.22)$$

gives the normalization condition (2.40).

Furthermore, elastic scattering requires that, as $Q^2 \rightarrow 0$,

$$(\eta - 1)q_- = \xi P_- \rightarrow \eta q_-, \quad (3.23)$$

and hence

$$\begin{aligned} \langle n | \mathcal{J}_+(P, P) | n \rangle \Big|_{e_1 \text{ term}} &= 2e_1 P_+ \frac{m_1^2}{M_n^2} \int_0^1 d\xi \frac{\Phi_n^2(\xi, M_0^2)}{\xi^2} \\ &= 2e_1 P_+ \mathcal{F}_n(0) = 2e_1 P_+, \end{aligned} \quad (3.24)$$

which follows from the wave function identity (B4). Note that *the charge is properly normalized and conserved for both components because we used the correct dressed quark current* (3.6).

Finally, we point out that the elastic transverse current is zero for all Q^2 , as required by the symmetries of the states

$$\langle n | \mathcal{J}_x(P_f, P) | n \rangle = 0. \quad (3.25)$$

We now turn to a discussion of deep inelastic scattering.

IV. DEEP INELASTIC SCATTERING

A. Extracting the structure functions from the DIS cross section

Using the standard conventions (also defined in Ref. [14]), the DIS cross section is

$$\begin{aligned} \frac{d^2\sigma}{d\Omega' dE'} &= \left(\frac{2\alpha E'}{Q^2} \right)^2 \left[W_2 \cos^2 \left(\frac{\theta}{2} \right) + 2W_1 \sin^2 \left(\frac{\theta}{2} \right) \right] \\ &= \sigma_M \left[W_2 + 2W_1 \tan^2 \left(\frac{\theta}{2} \right) \right], \end{aligned} \quad (4.1)$$

where the structure functions W_1 and W_2 are part of the hadronic tensor, defined by

$$\begin{aligned}\frac{W_{\mu\nu}}{4\pi M} &= -\left(g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2}\right) W_1 + \left(P_\mu - q_\mu \frac{P \cdot q}{q^2}\right) \left(P_\nu - q_\nu \frac{P \cdot q}{q^2}\right) \frac{W_2}{M^2} \\ &= -\left(g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2}\right) W_1 + \left(P_\mu + \frac{q_\mu}{2x}\right) \left(P_\nu + \frac{q_\nu}{2x}\right) \frac{W_2}{M^2},\end{aligned}\quad (4.2)$$

where $M = M_1$ is the target, or ground state mass. The structure function W_1 is immediately extracted from the xx (or yy) component of the tensor

$$MW_1 = \frac{W_{xx}}{4\pi} \rightarrow F_1(x) = \frac{1}{2}f(x). \quad (4.3)$$

Here we anticipate that MW_1 scales in the deep inelastic limit to the function $F_1(x)$ of the variable $x = Q^2/(2P \cdot q)$. Note that W_1 must be identically zero if the x (and y) components of the current are zero.

The structure function W_2 can be extracted from the $+$ and $-$ components of the current using

$$W_{-+} = \frac{1}{2}(W_{00} - W_{zz}) = 2\pi M \left\{ W_2 \left(1 + \frac{Q^2}{4x^2 M^2} \right) - W_1 \right\}. \quad (4.4)$$

In the deep inelastic limit this gives

$$\nu W_2 \rightarrow F_2(x) = 2xF_1(x) + \frac{1}{\pi}xW_{-+}. \quad (4.5)$$

The contribution from W_{-+} will be shown to go like ν^{-1} in the following sections. Hence, if F_1 is non-zero, we obtain the Callan-Gross relation. However, if the transverse components of the current are omitted, $F_1 = 0$ and νW_2 does not scale. In this case it is the quantity $\nu^2 W_2$ that scales, and we recover the results of Einhorn [3].

We now turn to a calculation of the structure functions. We calculate them first (i) in the *partonic picture*, where all final state interactions are ignored and the quarks in the final state are assumed to be free, and then (ii) in the *hadronic picture*, where confinement ensures that the only possible final states are the bound $q\bar{q}$ states we have already discussed. We emphasize that the partonic picture can never actually occur because the confining interaction, which also acts in the final state, can never be ignored.

B. DIS in the partonic picture

Consider deep inelastic scattering (DIS) in the partonic picture, which assumes a final state composed of free quarks with no interaction. This picture is illustrated in Fig. 11. The full result is the sum of a contributions in which the photon is absorbed by the quark with charge e_1 , and one in which it is absorbed by the antiquark with charge e_2 . In the center of mass of the ejectiles, either particle can go forwards or backwards and the unpolarized cross section is the sum over both helicities of the particles.

Begin by calculating the transverse component (chosen be be x) of the current. The outgoing quark has a mass of m_1 and momentum p_1 in the center of mass frame of the




FIG. 11: One of two diagrams that contribute to DIS in the partonic picture. Note the labeling of the momenta.

ejectiles, while the antiquark has a mass of m_2 and momentum of p_2 (in this section p_2 will be the physical momentum of the outgoing antiquark, opposite in sign to that used in the earlier sections of this paper). The current is the sum of two diagrams, only one of which is shown in Fig. 11. Since the $q\bar{q}$ vertex contains the factor of γ_- , only the factor $m + p_- \gamma_+$ in the numerator of the dressed propagator (2.30) survives, but we will retain only the factor of $p_- \gamma_+$ which dominates the result at large Q^2 . The following result is obtained for the x component of the current for the disintegration of the ground state (the subscript $n = 1$ usually attached to the ground state will be omitted in this section):

$$J_x = -e_1 [\bar{u}(p_1) \gamma_x \gamma_+ \gamma_- v(p_2)] \frac{(p_1 - q)_- G(p_1 - q, -p_2)}{m_1^2 - (p_1 - q)^2} + e_2 [\bar{u}(p_1) \gamma_- \gamma_+ \gamma_x v(p_2)] \frac{(p_2 - q)_- G(p_1, -p_2 + q)}{m_2^2 - (p_2 - q)^2}. \quad (4.6)$$

Introducing the momentum of the bound state, $P = p_1 + p_2 - q$, and defining

$$z = \frac{(p_1 - q)_-}{P_-} = \frac{(P - p_2)_-}{P_-} \quad (4.7)$$

[as in Eq. (2.36)], the first denominator can be written

$$\begin{aligned} m_1^2 - (p_1 - q)^2 &= m_1^2 - (P - p_2)^2 = m_1^2 - 2 \left[\frac{M^2}{2P_-} - \frac{m_2^2}{2p_{2-}} \right] (P - p_2)_- \\ &= m_1^2 - z \left(M^2 - \frac{m_2^2}{1 - z} \right) = z \left[\frac{m_1^2}{z} + \frac{m_2^2}{1 - z} - M^2 \right]. \end{aligned} \quad (4.8)$$

Here we have made use of the constraints $M^2 = 2P_- P_+$ and $m_2^2 = 2p_{2-} p_{2+}$. A similar result follows for the second denominator if we define a new momentum fraction z'

$$z' = \frac{(p_2 - q)_-}{P_-} = \frac{(P - p_1)_-}{P_-}. \quad (4.9)$$

With these definitions we obtain

$$J_x = -e_1 [\bar{u}(p_1) \gamma_x \gamma_+ \gamma_- v(p_2)] \frac{P_- G(z, P)}{\Delta(z, M^2)}$$

$$\begin{aligned}
& + e_2 [\bar{u}(p_1) \gamma_- \gamma_+ \gamma_x v(p_2)] \frac{P_- G(1 - z', P)}{\Delta(1 - z', M^2)} \\
& = -\frac{e_1}{\mathcal{N}} [\bar{u}(p_1) \gamma_x \gamma_+ \gamma_- v(p_2)] \Phi(z, P) \\
& \quad + \frac{e_2}{\mathcal{N}} [\bar{u}(p_1) \gamma_- \gamma_+ \gamma_x v(p_2)] \Phi(1 - z', P), \tag{4.10}
\end{aligned}$$

where the answer has been expressed in terms of the wave function (2.39).

The current and the DIS cross section are evaluated in the c.m. frame, using the kinematics defined in Appendix E. The unpolarized cross section is obtained by squaring the current and summing over spins using

$$\begin{aligned}
\sum_{\text{spins}} \bar{u}(p_1) \gamma_x \gamma_+ \gamma_- v(p_2) \bar{v}(p_2) \gamma_- \gamma_+ \gamma_x u(p_1) &= p_{1+} p_{2-} \text{trace} [\gamma_+ \gamma_- \gamma_+ \gamma_x \gamma_- \gamma_x \gamma_+ \gamma_-] \\
&= 32 p_{1+} p_{2-}, \tag{4.11}
\end{aligned}$$

and neglecting the interference term (which vanishes in the large Q^2 limit). Observing that $z \rightarrow z' \rightarrow x$ in the DIS limit, and summing over all colors of the outgoing quarks (which do not interfere) gives

$$\begin{aligned}
\sum_{\text{spins}} |J_x|^2 &= \frac{32 N_c}{\mathcal{N}^2} \{ e_1^2 p_{1+}^{(1)} p_{2-}^{(1)} \Phi^2(x, P) + e_2^2 p_{1-}^{(2)} p_{2+}^{(2)} \Phi^2(1 - x, P) \} \\
&= 4\pi Q^2 \left(\frac{1 - x}{x} \right) \{ e_1^2 \Phi^2(x, P) + e_2^2 \Phi^2(1 - x, P) \}, \tag{4.12}
\end{aligned}$$

where the momentum are evaluated in Appendix E. The hadronic tensor defined in Eq. (4.2) is related to the square of the currents by

$$\frac{W_{\mu\nu}}{4\pi M} = \int \frac{dp_1 dp_2}{(2\pi)^2 4p_{10} p_{20}} (2\pi)^2 \delta^2(P_f - P - q) \sum_{\text{spins}} J_\mu J_\nu, \tag{4.13}$$

The W_{xx} component of the tensor is [14] is therefore

$$\begin{aligned}
W_{xx} &= 4\pi M W_1 = \frac{1}{4 |p_z| (p_{10} + p_{20})} \sum_{\text{spins}} |J_x|^2 \\
&= 2\pi \{ e_1^2 \Phi^2(x, P) + e_2^2 \Phi^2(1 - x, P) \}. \tag{4.14}
\end{aligned}$$

This gives the familiar parton model result for the structure function W_1

$$M W_1 = F_1(x) = \frac{1}{2} \{ e_1^2 \Phi^2(x, P) + e_2^2 \Phi^2(1 - x, P) \} = \frac{1}{2} f(x). \tag{4.15}$$

The structure function W_2 , given in Eq. (4.5), is then

$$\nu W_2 = 2MxW_1 = 2xF_1(x) = F_2(x) = xf(x), \tag{4.16}$$

where we anticipate the result $W_{-+} \rightarrow 0$ as $Q^2 \rightarrow \infty$. This is the Callan-Gross relation.

Now look at the plus and minus components of the current. If the Dirac space were restricted to two dimensions, only these components of the current would exist. The minus component can be obtained immediately from (4.10) by replacing γ_x with γ_- , giving

$$J_- = -[\bar{u}(p_1) \gamma_- v(p_2)] \frac{2}{\mathcal{N}} \{ e_1 \Phi(z, P) - e_2 \Phi(1 - z', P) \}, \tag{4.17}$$

The plus component could include the two contributions from the quark current, Eq. (3.6), but in the partonic picture the outgoing quark is not dressed, so the second part of the current proportional to g^2 should be ignored. However, in order to study the contribution of this term we will retain it for now. The leading contribution from both terms in the J_+ current is

$$J_+ = -\frac{e_1}{\mathcal{N}} \left\{ m_{01} \frac{\Phi(z, P)}{p_{1-}^{(1)} - q_-} [\bar{u}(p_1) \gamma_+ \gamma_- v(p_2)] + \frac{g^2}{\pi} \frac{\Phi(z, P)}{p_{1-}^{(1)}(p_{1-}^{(1)} - q_-)} [\bar{u}(p_1) \gamma_- v(p_2)] \right\} \\ + \frac{e_2}{\mathcal{N}} \left\{ m_{02} \frac{\Phi(1 - z', P)}{p_{2-}^{(2)} - q_-} [\bar{u}(p_1) \gamma_- \gamma_+ v(p_2)] + \frac{g^2}{\pi} \frac{\Phi(z, P)}{p_{2-}^{(2)}(p_{2-}^{(2)} - q_-)} [\bar{u}(p_1) \gamma_- v(p_2)] \right\}. \quad (4.18)$$

The W_{-+} component of the hadron tensor is then

$$W_{-+} = \frac{1}{4 |p_z| (p_{10} + p_{20})} \sum_{\text{spins}} J_- J_+ = \frac{x}{2Q^2(1-x)} \sum_{\text{spins}} J_- J_+. \quad (4.19)$$

Using (4.17) and (4.18), and summing over colors, the e_1^2 contribution to the current sum becomes, at large Q^2 ,

$$\sum_{\text{spins}} J_- J_+ |_{e_1 \text{ term}} = \frac{32 e_1^2 N_c}{\mathcal{N}^2} \Phi^2(x, P) \left[\frac{p_{2-}^{(1)}}{p_{1-}^{(1)} - q_-} \right] \left(m_{01}^2 - \frac{g^2}{\pi} \right) \\ = 8\pi e_1^2 m_1^2 \left(\frac{1-x}{x} \right) \Phi^2(x, P). \quad (4.20)$$

Hence

$$W_{-+} |_{e_1 \text{ term}} = \frac{4\pi e_1^2}{Q^2} m_1^2 \Phi^2(x, P) = 4\pi \frac{m_1^2}{Q^2} f(x) |_{e_1 \text{ term}}, \quad (4.21)$$

and this term is subleading at high Q^2 . Note that the effect of neglecting the term proportional to g^2 in the $+$ component of the quark current (as we are instructed to do in the partonic picture) is to replace m_1^2 by the bare quark mass m_{01}^2 . If $W_1 = 0$, then $\nu^2 W_2$ scales as

$$\nu^2 W_2 |_{e_1 \text{ term}} \rightarrow \frac{Q^2}{2\pi M} W_{-+} |_{e_1 \text{ term}} = 2 \frac{m_{01}^2}{M} f(x) |_{e_1 \text{ term}}. \quad (4.22)$$

Up to a factor, this is the result originally obtained by Einhorn [3]. In that paper, the Dirac space was restricted to 2 dimensions, W_1 was necessarily zero, and $\nu^2 W_2$ scaled as (4.22). This distribution amplitude does not have the correct physical interpretation. In particular, it predicts that the DIS cross section depends on the square of bare quark mass, a clearly unphysical result. Only by considering the full four dimensional Dirac space are we able to obtain scaling for νW_2 , and the familiar physical result (4.15) for $f(x)$.

Another consequence of the fact that W_{-+} is subleading is that we can avoid coming to grips with the fact that the current components J_- and J_+ , as defined above, do not conserve current. Since these components do not contribute to the final result, we need not discuss how they can be redefined in order to conserve current.

C. DIS in the hadronic picture

1. Exact results for inelastic scattering

The *exact* hadronic tensor (4.2) for electroproduction of a *single* final state f with mass M_n (denoted by n) from the ground state i with mass $M_1 \equiv M$ (denoted by 0) is

$$\begin{aligned} W_{\mu\nu}|_f &= \int \frac{dP_{fz}}{(2\pi)2P_{f0}} (2\pi)^2 \delta^2(P_f - P - q) \sum_{\text{spins}} \langle 0|\mathcal{J}_\mu|n\rangle \langle n|\mathcal{J}_\nu|0\rangle \\ &= \delta(M_n - P_0 - q_0) \frac{\pi}{M_n} \sum_{\text{spins}} \langle 0|\mathcal{J}_\mu|n\rangle \langle n|\mathcal{J}_\nu|0\rangle . \end{aligned} \quad (4.23)$$

This is a delta function. In any physical measurement, the detectors will accept a finite range of values of the final electron energy, E' , requiring that this theoretical cross section be averaged over E' . This averaging process will be discussed in subsection IV D below, where the delta function will be written as a delta function in the Bjorken variable x , which spikes at values of x_n corresponding to the excitation of the final state n . In the c.m. system, this leads to the following *exact* transformation

$$\delta(M_n - P_0 - q_0) = \delta(x - x_n) \frac{2x_n^2 M_n}{Q^2} \quad (4.24)$$

with x_n the value of x at the bound state mass M_n , given in Eq. (4.26) below.

In the DIS limit, the transverse currents dominate the scattering. The exact result for the transverse tensor, W_{xx} , that describes the scattering from the ground state with γ_- structure, to a *single* transverse final state n with $\gamma_- \gamma_x$ structure, is

$$W_{xx}|_n = \delta(x - x_n) \frac{2\pi x_n^2}{Q^2} \left\{ \mathcal{J}_n^{1,x}(Q^2, y) + \mathcal{J}_n^{2,x}(Q^2, y) \right\}^2, \quad (4.25)$$

where the reduced transverse current was defined in Eq. (3.18). (Here the *fi* subscript is replaced by n , and denotes a transition between the ground and n th state.) Next we observe that the value of x at which the final state is excited is

$$x_n = \frac{Q^2}{M_n^2 - M^2 + Q^2}, \quad (4.26)$$

and that the spacing between neighboring levels is

$$\Delta_{x_n} = x_n - x_{n+1} = \frac{\delta M_n^2}{Q^2} x_n x_{n+1}, \quad (4.27)$$

with $\delta M_n^2 = M_{n+1}^2 - M_n^2$. Hence the *exact* result for the total inelastic tensor W_{xx} is

$$W_{xx} = \sum_n \delta(x - x_n) \Delta_{x_n} \left(\frac{2\pi}{\delta M_n^2} \right) \left(\frac{x_n}{x_{n+1}} \right) \left\{ \mathcal{J}_n^{1,x}(Q^2, y) + \mathcal{J}_n^{2,x}(Q^2, y) \right\}^2, \quad (4.28)$$

where the sum is over all possible final states n .

2. Hadronic result in the DIS limit

The DIS limit is defined by the requirement that $Q^2 \rightarrow \infty$ with x held constant. In this limit both Q^2 and M_n^2 become large, with

$$x_n \rightarrow \frac{Q^2}{M_n^2 + Q^2}, \quad (\text{as } Q^2 \text{ and } M_n^2 \rightarrow \infty). \quad (4.29)$$

Study of the DIS limit therefore requires estimating the transition form factors for large final state number n , but for all x (since x_n depends on Q^2). In the DIS limit, $y \rightarrow x$ [easily seen from the expansions (E1)], so y will be replaced by x in the following discussion.

When n is large the “tails” of the wave functions are very small, and the exact transition current (3.18) can be approximated by neglecting the contribution from the region $\xi \in [0, x]$ (where $\xi' < 0$), giving

$$\begin{aligned} \mathcal{J}_n^{1,x}(Q^2, x) &\simeq \frac{e_1 m_{01} x}{1-x} \int_y^1 d\xi \frac{\Phi_n(\xi', P_f) \Phi(\xi, P)}{\xi' \xi} F_1^{\text{eff}}(\eta, Q^2) \\ &= e_1 m_{01} x \int_0^1 d\xi' \frac{\Phi_n(\xi', P_f) \Phi([\xi'(1-x) + x], P)}{\xi' [\xi'(1-x) + x]} F_1^{\text{eff}}(\eta, Q^2), \end{aligned} \quad (4.30)$$

where we continue to suppress the label $n = 1$ on the ground (initial) state wave function, and

$$\eta = -\xi' \left(\frac{1-x}{x} \right) < 0 \quad (4.31)$$

in the region of integration. When $\eta < 0$ and $Q^2 \rightarrow \infty$, the relation (D10) may be used to approximate the quark form factor by

$$F_i(\eta, Q^2) \rightarrow -\frac{g^2}{\pi Q^2 \eta (1-\eta)}, \quad (4.32)$$

giving

$$F_1^{\text{eff}}(\eta, Q^2) \rightarrow 1. \quad (4.33)$$

Equation (4.30) is now further reduced by expanding $\Phi(\xi, P)/\xi$ around $\xi = x$

$$\frac{\Phi(\xi, P)}{\xi} = \frac{\Phi(x, P)}{x} + \delta \frac{d}{dx} \left(\frac{\Phi(x, P)}{x} \right) + \frac{1}{2} \delta^2 \frac{d^2}{dx^2} \left(\frac{\Phi(x, P)}{x} \right) + \dots \quad (4.34)$$

where the higher order terms are proportional to powers of δ

$$\delta \equiv \xi'(1-x). \quad (4.35)$$

This replacement gives the following series for $\mathcal{J}$

$$\begin{aligned} \mathcal{J}_n^{1,x}(Q^2, x) &= e_1 m_{01} \Phi(x, P) \int_0^1 d\xi' \frac{\Phi_n(\xi', P_f)}{\xi'} \\ &\quad + e_1 m_{01} \sum_{m=1}^{m=\infty} d_n^m \frac{(1-x)^m}{m!} \left(\frac{d}{dx} \right)^m \left(\frac{\Phi(x, P)}{x} \right) \end{aligned} \quad (4.36)$$

where

$$d_n^m = \int_0^1 d\xi' \xi'^{m-1} \Phi_n(\xi', P_f). \quad (4.37)$$

Using the identity (B10) and the other estimates given in Appendix B it follows that

$$\lim_{n \rightarrow \infty} \frac{d_n^m}{C_n} \rightarrow \frac{\text{const}}{n} \rightarrow 0. \quad (4.38)$$

Hence the large n approximation to (4.36) is

$$\begin{aligned} \mathcal{J}_n^{1,x}(Q^2, x) &\rightarrow e_1 m_{01} \Phi(x, P) \int_0^1 d\xi' \frac{\Phi_n(\xi', P_f)}{\xi'} \\ &= e_1 C_n \Phi(x, P). \end{aligned} \quad (4.39)$$

The plus component of the transition current is smaller than the transverse current by a factor of m_{01}/q_- [compare the two results in Eq. (3.16)], and hence is obtained immediately from the result (4.39). Using Eq. (E1) for q_- , and Eq. (4.29) to replace $\sqrt{(1-x_n)/x_n}$, gives

$$\mathcal{J}_n^{1,+}(Q^2, x) \simeq -\sqrt{2} e_1 C_n \frac{m_{01} M_n}{Q^2} \Phi(x, P). \quad (4.40)$$

The minus component can also be obtained from Eq. (4.39). Reviewing the derivation, we see that the minus component, in the DIS limit, is obtained from the transverse component by replacing

$$r \equiv \frac{m_{01}}{2\eta(1-\eta)q_-} = \frac{m_{01}y}{\sqrt{2}\xi'\xi M_n} \rightarrow 1 \quad (4.41)$$

Hence, multiplying (4.39) by r^{-1}

$$\begin{aligned} \mathcal{J}_n^{1,-}(Q^2, x) &\simeq \sqrt{2} e_1 \Phi(x, P) \left\{ M_n \int_0^1 d\xi^* \Phi_f(\xi^*, P_f) \right\} \\ &\simeq \sqrt{2} e_1 C_n \left(\frac{m_{01} + (-1)^{n-1} m_{02}}{M_n} \right) \Phi(x, P), \end{aligned} \quad (4.42)$$

where the ξ' integral was evaluated using identity (B11).

The transition currents can be shown to be gauge invariant by using the Ward-Takahashi identity on the quark-photon vertex, and reducing the result using the Bethe-Salpeter equation.

These currents will now be used to determine the structure functions W_1 and W_2 in the hadron picture.

3. The DIS cross section from the transition form factors

The cross section in the DIS limit can now be obtained from the exact formula (4.28). As long as x is not too close to unity each term in the sum must correspond to some state with large n (the states with small n “pile up” near $x = 1$ where their total contribution

is very small), and the approximate result for the transverse current, (4.39), may be used. Furthermore, for large n , δM_n^2 may be approximated by using (B13), and $x_{n+1} \simeq x_n$. Hence the final hadronic result [denoted $\tilde{f}$ to distinguish it from the partonic result f of Eq. (4.15)] in the DIS limit is

$$\tilde{f}(x) = 2\tilde{F}_1(x) = \frac{W_{xx}}{2\pi} = \sum_n \delta(x - x_n) \Delta_{x_n} \left\{ e_1 \Phi(x, P) + (-1)^n e_2 \Phi(1 - x, P) \right\}^2. \quad (4.43)$$

Note that $\tilde{F}_1(x)$ is a series of separate spikes, zero for all x except at particular values x_n , where it is infinite. The partonic function, $F_1(x)$, is a smooth function of x . To compare $\tilde{F}_1(x)$ to $F_1(x)$, we must average over x . This will lead to the concept of duality and will be discussed in the next subsection below.

Now look at the W_{-+} component of the tensor. Replacing the transverse components of the current in Eq. (4.28) with plus and minus components, and taking the DIS limit, gives

$$W_{-+} = \frac{2\pi}{C_\infty^2} \sum_n \delta(x - x_n) \Delta_{x_n} \times \left\{ \mathcal{J}_n^{1,-}(Q^2, x) + \mathcal{J}_n^{2,-}(Q^2, x) \right\} \left\{ \mathcal{J}_n^{1,+}(Q^2, x) + \mathcal{J}_n^{2,+}(Q^2, x) \right\}^*. \quad (4.44)$$

Substituting Eqs. (4.40) and (4.42), and dropping the term proportional to $(-1)^{n-1}$ (which averages to zero) gives

$$W_{-+}|_{e_1 \text{ term}} = \frac{4\pi m_{01}^2}{Q^2} \sum_n \delta(x - x_n) \Delta_{x_n} e_1^2 \Phi^2(x, P). \quad (4.45)$$

This is a nonleading term and can be neglected in the DIS limit. Hence the hadronic picture also gives the Callan-Gross relation, and it is sufficient to compare the functions F_1 and $\tilde{F}_1$ only.

If W_1 were zero, Eq. (4.45) would give the following scaling relation for $\nu^2 W_2$

$$\nu^2 W_2|_{e_1 \text{ term}} \rightarrow \frac{Q^2}{2\pi M} W_{-+}|_{e_1 \text{ term}} = 2 \frac{m_{01}^2}{M} \tilde{f}(x)|_{e_1 \text{ term}}. \quad (4.46)$$

This is to be compared with (4.22). Hence, the duality of W_1 and the duality of the non-leading terms in $\nu^2 W_2$ depends on comparison of the same functions, $f(x)$ and $\tilde{f}(x)$.

We discuss this comparison now.

D. Duality and its implications

In any physical measurement, the detectors which define the final state will accept a range of final electron energies $\delta E'$. For fixed Q^2 , this can be converted into the acceptance of a range of values of x centered at x_i with width δx so that x varies over the interval bounded by $x_{i\pm} = x_i \pm \delta x/2$. Then the *experimentally measured* hadronic structure function in the DIS limit can be computed from (4.43)

$$\begin{aligned} \langle \tilde{f} \rangle_{x_i} &\equiv \frac{1}{\delta x} \int_{x_{i-}}^{x_{i+}} dx \tilde{f}(x) \\ &= \frac{1}{\delta x} \sum_{n \in x_i} \Delta_{x_n} \left| e_1 \Phi(x_n, P) + (-1)^{n-1} e_2 \Phi(1 - x_n, P) \right|^2. \end{aligned} \quad (4.47)$$




FIG. 12: Quark-hadron duality

Note that this is the sum of smooth terms proportional to e_1^2 and e_2^2 , and a rapidly oscillating interference term proportional to $e_1 e_2$. In the DIS limit, the separation Δ_x between states approaches zero, and therefore a large number of states are necessarily included in any interval δx . The interference term, which changes sign as $(-1)^{n-1}$, therefore averages to zero. This cancellation of the interference term was pointed out originally by Einhorn [3], and emphasized recently by Close and Isgur in the context of the nonrelativistic quark model [19].

The smooth terms can be approximated by their value at the center of the interval. Using the fact that the number of states in the interval times the x spacing between them must necessarily equal the width δx , so that

$$\sum_{n \in x_i} \Delta_{x_n} = \delta x. \quad (4.48)$$

we can reduce (4.47) to

$$\begin{aligned} \langle \tilde{f} \rangle_{x_i} &\simeq \left\{ e_1^2 \Phi^2(x_i, P) + e_2^2 \Phi^2(1 - x_i, P) \right\} \frac{1}{\delta x} \sum_{n \in x_i} \Delta_{x_n} \\ &\quad + 2e_1 e_2 \frac{1}{\delta x} \sum_{n \in x_i} (-1)^{n-1} \Delta_{x_n} \Phi(x_n, P) \Phi(1 - x_n, P) \\ &\rightarrow e_1^2 \Phi^2(x_i, P) + e_2^2 \Phi^2(1 - x_i, P) = f(x_i), \end{aligned} \quad (4.49)$$

where $f(x)$ was defined in Eq. (4.15).

This result should be compared with the partonic result, averaged in the same way. Since the partonic result is smooth (and the interference term already neglected) the partonic average is trivial

$$\langle f \rangle_{x_i} \equiv \frac{1}{\delta x} \int_{x_{i-}}^{x_{i+}} dx f(x) \simeq f(x_i) \frac{1}{\delta x} \int_{x_{i-}}^{x_{i+}} dx = f(x_i). \quad (4.50)$$

Hence we see that

$$\langle \tilde{f} \rangle_{x_i} = \langle f \rangle_{x_i}, \quad (4.51)$$

proving duality.

Note that a similar identity holds for the n th moments of f . In the high Q^2 limit when the final states are numerous and close enough together so that $\Delta_{x_n} \rightarrow dx \rightarrow 0$, the m th moment of the hadronic structure function is

$$\begin{aligned} \langle x^m \rangle &\equiv \int_0^1 dx x^m \tilde{f}(x) = \sum_n \Delta_{x_n} x_n^m \left[e_1^2 \Phi^2(x_n, P) + e_2^2 \Phi^2(1 - x_n, P) \right] \\ &\rightarrow \int_0^1 dx x^m \left[e_1^2 \Phi^2(x, P) + e_2^2 \Phi^2(1 - x, P) \right] \\ &= \int_0^1 dx x^m f(x). \end{aligned} \tag{4.52}$$

Quark-hadron duality means that properly averaged hadronic observables in a certain kinematic regime (high Q^2) can be described by perturbative QCD, as schematically represented in Fig. 12 and demonstrated in Eq. (4.51). Recent data from Jefferson Laboratory exhibit duality, and this has focused attention on this old subject. Recent studies show that the weak decays within the 't Hooft model exhibit duality [11, 12, 13], and in the framework of DIS, Close and Isgur [19] studied its emergence in the context of realistic nonrelativistic quark models. A recent paper by Isgur, Jeschonnek, Melnitchouk and Van Orden [20] studied duality using a relativistic Klein-Gordon equation with a confining interaction. This work is complementary to these other studies.

While we have proved that duality emerges in the DIS limit, and have corrected the pathologies of earlier work, we have not yet studied how duality emerges as a function of Q^2 . A future paper will present numerical studies of the onset of duality for rQCD [15].

V. CONCLUSIONS

We have shown that a very satisfactory description of deep inelastic scattering can be obtained from large N_c QCD in 1+1 dimensions, *provided* the quark spin degrees of freedom are extended to the full 1+3 dimensions (along with the leptons and the electromagnetic field). We refer to this new model as reduced QCD, and suggest that rQCD may be used to provide insight into the behavior of many other physical processes. In the applications to DIS discussed in this paper, we are able to confirm the Callan-Gross relation, and that duality emerges in the deep inelastic limit.

Our theoretical discussion teaches several lessons. Electromagnetic gauge invariance and duality emerge only if the quark current is fully and consistently dressed. The way in which a quark form factor emerges from this dynamical model mimics the phenomenological approach of Gross and Riska [18], showing that their prescription has a dynamical justification. Our treatment of confinement removes all singularities, giving a finite mass for the dressed quark. This mass is gauge dependent, showing that it is not a physical quantity, which can be taken as a demonstration of confinement. But the fact that we have finite dressed quark masses in the presence of confinement provides justification for the constituent quark model. The simultaneous presence of confinement and finite quark mass is possible only if the “on-shell” quark scattering amplitude is identically zero, and we have shown that this is indeed true, as modeled previously in Ref. [17].

The treatment of reduced QCD presented here lays a foundation for much further study.

VI. ACKNOWLEDGMENTS

This work has been supported in part by the DOE through grant No. DE-FG05-88ER40435. The Southeastern Universities Research Association (SURA) operates the Thomas Jefferson National Accelerator Facility under DOE contract DE-ACO5-84er40150. One of us (Z. B.) has been supported by a Predoctoral Fellowship at the College of William & Mary and by the Fundação para a Ciência e a Tecnologia under the post doctoral grant SFRH/BPD/5661/2001 the project grant CFIF-Plurianual.

APPENDIX A: REDUCTION OF THE TWO QUARK INTERACTION

Integrals over the variable k_+ like the one encountered in Eq. (2.35) appear several places in this paper. Here we evaluate it in full generality.

Consider the integral

$$I_2(p_-, r) = 4i \int \frac{d^2k}{(2\pi)^2} \frac{V(p, k) k_- (k_- - r_-) F(k_-, r)}{[m_1^2 - k^2 - i\epsilon] [m_2^2 - (k - r)^2 - i\epsilon]}. \quad (\text{A1})$$

Substituting for V gives

$$I_2(p_-, r) = \frac{ig^2}{\pi^2} \int \frac{d^2k}{(p_- - k_-)^2} \left\{ \frac{F(k_-, r)}{d_2(k_-, k_+, r)} - \frac{F(p_-, r)}{d_2(p_-, k_+, r)} \right\}, \quad (\text{A2})$$

where the denominator d_2 is

$$\begin{aligned} d_2(k_-, k_+, r) &= \left[\frac{m_1^2}{k_-} - 2k_+ - i \frac{\epsilon}{k_-} \right] \left[\frac{m_2^2}{(k - r)_-} - 2(k - r)_+ - i \frac{\epsilon}{(k - r)_-} \right] \\ &= \left[\frac{m_1^2}{y r_-} - 2k_+ - i \frac{\epsilon}{y r_-} \right] \left[-\frac{m_2^2}{(1 - y) r_-} - 2(k - r)_+ + i \frac{\epsilon}{(1 - y) r_-} \right], \end{aligned} \quad (\text{A3})$$

and we have introduced the momentum fractions (2.36). This shows that the denominator has only two poles in k_+ , and that they will both be in the same half of the complex plane (giving zero for the integral) unless

$$0 \leq y \leq 1. \quad (\text{A4})$$

Doing the integral over k_+ , and expressing the answer in terms of the momentum fractions y and z gives

$$\begin{aligned} I_2(z r_-, r) \rightarrow I_2(z, r) &= \frac{g^2}{\pi} \int_0^1 \frac{dy}{(z - y)^2} \left\{ \frac{F(y r_-, r)}{\Delta(y, r^2)} - \theta(z) \theta(1 - z) \frac{F(z r_-, r)}{\Delta(z, r^2)} \right\} \\ &\quad - \theta(z) \theta(1 - z) \frac{g^2 F(z r_-, r)}{\pi \Delta(z, r^2)} \left\{ \int_{-\infty}^0 + \int_1^{\infty} \right\} \frac{dy}{(z - y)^2}, \end{aligned} \quad (\text{A5})$$

where the new denominator $\Delta(y, r^2)$ is

$$\Delta(y, r^2) = \left[\frac{m_1^2}{y} + \frac{m_2^2}{(1 - y)} - r^2 \right]. \quad (\text{A6})$$

[Note that this result holds for *both* signs of r_- . This is because the sign of the contour integral, which depends on the sign of r_- , is cancelled by the sign of the integral over $k_- = yr_-$, which must be changed from $\int_1^0 \rightarrow -\int_0^1$ when $r_- < 0$.] It is important to note that the integral over y is not restricted in the term in (A5) that is proportional to $F(zr_-, r)$ because for that term the restriction (A4) applies to z and not y . This term not only regulates the singularity in the first term (which arises only if $0 \leq z \leq 1$), but also contributes an additional contribution. Furthermore, if z lies outside of the interval $0 \leq z \leq 1$, the first term is non singular (except at the end points of the interval) and is nonzero! Hence, in the region $0 \leq z \leq 1$ the integral (A5) becomes

$$\begin{aligned} I_2(z, r) &= \frac{g^2}{\pi} \int_0^1 \frac{dy}{(z-y)^2} \left\{ \frac{F(yr_-, r)}{\Delta(y, r^2)} - \frac{F(zr_-, r)}{\Delta(z, r^2)} \right\} - \frac{g^2 F(zr_-, r)}{\pi z(1-z) \Delta(z, r^2)} \\ &= \int_0^1 dy V(z, y) \frac{F(yr_-, r)}{\Delta(y, r^2)} - \frac{g^2/\pi}{z(1-z)} \frac{F(zr_-, r)}{\Delta(z, r^2)} \quad (\text{if } z \in [0, 1]) , \end{aligned} \quad (\text{A7})$$

and outside of this region it is

$$I_2(z, r) = \frac{g^2}{\pi} \int_0^1 \frac{dy}{(z-y)^2} \frac{F(yr_-, r)}{\Delta(y, r^2)} = \int_0^1 dy V_0(z, y) \frac{F(yr_-, r)}{\Delta(y, r^2)} \quad (\text{if } z \notin [0, 1]) , \quad (\text{A8})$$

where we have defined

$$\frac{(r_-)^2}{\pi} V(p, k) \rightarrow \begin{cases} V(z, y) \equiv \frac{g^2}{\pi} \left\{ \frac{1}{(z-y)^2} - \delta(z-y) \int_0^1 dy' \frac{1}{(z-y')^2} \right\} & \text{if } z \in [0, 1] \\ V_0(z, y) \equiv \frac{g^2}{\pi} \frac{1}{(z-y)^2} & \text{if } z \notin [0, 1] \end{cases} . \quad (\text{A9})$$

The multiplication by r_-^2 in (A9) is needed to scale the momenta from $k_- \rightarrow y$, etc.

We emphasize that the identities (A7) and (A8) hold for both $r_- > 0$ and $r_- < 0$. The first case is needed for the bound state equation and the second for electron scattering.

APPENDIX B: IDENTITIES INVOLVING BOUND STATE WAVE FUNCTIONS

In this appendix we derive a number of identities that hold when $\Phi_n(0, r) = \Phi_n(1, r) = 0$ (i.e. in all cases but the chiral limit). Equation (2.42) leads immediately to the following identity

$$\int_0^1 dz \left(\frac{m_1^2}{z} + \frac{m_2^2}{1-z} \right) \Phi'_n(z, r) \Phi_n(z, r) = 0 . \quad (\text{B1})$$

Integrating by parts gives

$$m_1^2 \int_0^1 dz \frac{\Phi_n^2(z, r)}{z^2} = m_2^2 \int_0^1 dz \frac{\Phi_n^2(z, r)}{(1-z)^2} . \quad (\text{B2})$$

An additional identity is derived by multiplying (2.42) by $2(1-z) \Phi'_n(z, r)$ and integrating. First observe that

$$\begin{aligned} &2 \int_0^1 dz (1-z) \Phi'_n(z, r) \int_0^1 dy \frac{\Phi'_n(y, r)}{(z-y)} \\ &= \int_0^1 \int_0^1 dz dy \Phi'_n(z, r) \Phi'_n(y, r) \frac{y-z}{z-y} = 0 . \end{aligned} \quad (\text{B3})$$

Hence, integrating by parts and using the normalization condition (2.40) for the wave function, we obtain

$$M_n^2 = m_1^2 \int_0^1 dz \frac{\Phi_n^2(z, r)}{z^2}. \quad (\text{B4})$$

Similarly we can prove that

$$M_n^2 = m_2^2 \int_0^1 dz \frac{\Phi_n^2(z, r)}{(1-z)^2}. \quad (\text{B5})$$

Additional identities are needed for the study of duality. First, integrate Eq. (2.37) over z to obtain

$$M_n^2 \int_0^1 dz \Phi_n(z, r) = \int_0^1 dz \left(\frac{m_{01}^2}{z} + \frac{m_{02}^2}{1-z} \right) \Phi_n(z, r). \quad (\text{B6})$$

Next, the commutator of the operator

$$\mathcal{K} \Phi_n(x, r) \equiv \oint_0^1 dy \frac{\Phi_n(y, r)}{y-x} \quad (\text{B7})$$

with the Hamiltonian defined in (2.37) is

$$[H, \mathcal{K}] \Phi_n(x, r) = \int_0^1 dy \left[\frac{m_{01}^2}{xy} - \frac{m_{02}^2}{(1-x)(1-y)} \right] \Phi_n(y, r), \quad (\text{B8})$$

as discussed in Refs. [2, 7]. Multiplying this by $\Phi_n(x, r)$ and integrating over x gives zero on the l.h.s., and hence the relation

$$m_{01}^2 \left(\int_0^1 dz \frac{\Phi_n(z, r)}{z} \right)^2 = m_{02}^2 \left(\int_0^1 dz \frac{\Phi_n(z, r)}{1-z} \right)^2. \quad (\text{B9})$$

Taking the square root of both sides of this equation gives

$$m_{01} \int_0^1 dz \frac{\Phi_n(z, r)}{z} = \pm m_{02} \int_0^1 dz \frac{\Phi_n(z, r)}{1-z} \equiv C_n, \quad (\text{B10})$$

where the sign is positive if the phase of $\Phi_n(0, r)$ is the same as $\Phi_n(1, r)$ and negative if it is opposite. Since the states are non-degenerate, the phase of the n th eigenstate is $(-1)^{(n-1)}$. Combining this with Eq. (B6) gives

$$M_n^2 \int_0^1 dz \Phi_n(z, r) = [m_{01} + (-1)^{(n-1)} m_{02}] C_n, \quad (\text{B11})$$

where the constant C_n is yet to be determined. In this paper we have chosen the sign of $\Phi_n(z, r)$ so that it is always positive as $z \rightarrow 0$, so that $C_n > 0$ for all n .

Our demonstration of duality depends upon the relations

$$\lim_{n \rightarrow \infty} C_n \rightarrow C_\infty = g\sqrt{\pi} \quad (\text{B12})$$

$$\lim_{n \rightarrow \infty} (M_{n+1}^2 - M_n^2) \rightarrow C_\infty^2 = \pi g^2, \quad (\text{B13})$$

which can be derived approximately using arguments given by 't Hooft [1], and Callan, Coote, and Gross [2], given here for completeness. First, for large n , the boundary conditions require that the normalized wave function go like

$$\Phi_n(z, r) \sim \sqrt{2} \sin(n\pi z). \quad (\text{B14})$$

One test of this approximation is shown in Fig. 13. If n is very large, so that $M_n \gg m_{0i}$, the behavior of the wave function at the endpoints can be ignored, and near $z = 1/2$ Eq. (2.42) can be approximated

$$M_n^2 \Phi_n(z, r) \simeq -\frac{g^2}{\pi} \oint_{-\infty}^{\infty} \frac{\Phi'_n(y, r)}{y - z}. \quad (\text{B15})$$

Substituting the ansatz (B14) into this equation gives

$$\begin{aligned} M_n^2 \sin(n\pi z) &= -g^2 n \oint_{-\infty}^{\infty} \frac{\cos(n\pi y)}{y - z} = -g^2 n \operatorname{Re} \int_{-\infty}^{\infty} \frac{\cos(n\pi y)}{y - z - i\epsilon} \\ &= \pi g^2 n \sin(\pi n z). \end{aligned} \quad (\text{B16})$$

Hence $M_n^2 \rightarrow \pi g^2 n$ as $n \rightarrow \infty$ and identity (B13) is proved. To prove (B12) we consider the matrix element of (B8)

$$\begin{aligned} \int_0^1 \Phi_{n+1}(x, r) [H, \mathcal{K}] \Phi_n(x, r) &= 2C_{n+1}C_n \\ &= (M_{n+1}^2 - M_n^2) \oint_0^1 dx dy \frac{\Phi_{n+1}(x, r) \Phi_n(y, r)}{y - x}. \end{aligned} \quad (\text{B17})$$

where we used the fact that the phases of Φ_n and Φ_{n+1} are opposite. The integral is evaluated in the large n approximation

$$\begin{aligned} \oint_0^1 dx dy \frac{\Phi_{n+1}(x, r) \Phi_n(y, r)}{y - x} &\simeq 2 \operatorname{Re} \int_0^1 dx \int_{-\infty}^{\infty} dy \frac{\sin([n+1]\pi x) \sin(n\pi y)}{y - x - i\epsilon} \\ &= 2 \int_0^1 dx \sin([n+1]\pi x) \cos(n\pi x) \\ &= 2 \left(1 + \frac{1}{2n+1} \right) \rightarrow 2. \end{aligned} \quad (\text{B18})$$

Hence,

$$C_{n+1}C_n \simeq \pi g^2 \simeq C_\infty^2 \quad (\text{B19})$$

and (B12) is proved.

A numerical demonstration of the identities (B10), (B12), and (B13) is given in Figs. 13. The identities are satisfied to better than a few percent for state numbers between 40 and 80, provided the number of splines is at least 160 (twice the number of the maximum state of interest).

Two additional relations that follow from the completeness relation (2.51) and the bound state equation are needed. If $z \in [0, 1]$, then:

$$\sum_n \Phi_n(z, r_n) \int_0^1 dz' \Phi_n(z', r_n) = 1 \quad (\text{B20})$$

$$\sum_n \left(\frac{m_{01}^2}{z} + \frac{m_{02}^2}{1-z} - M_n^2 \right) \Phi_n(z, r_n) \int_0^1 dz' \Phi_n(z', r_n) = 0. \quad (\text{B21})$$




FIG. 13: Numerical test of the relations (B10), (B12), and (B13) for $m_{01} = 1.5$ and $m_{02} = 3$ and different numbers of splines, n_s . Solid circles are values of the l.h.s. of (B10) divided by C_∞ ; open circles are the absolute value of the r.h.s. divided by C_∞ . They should be equal for each state, and approach unity as $n \rightarrow \infty$. The tiny solid squares are the values of $(\mu_{n+1}^2 - \mu_n^2)/\pi^2$ which, according (B13), approach unity as $n \rightarrow \infty$. The open triangles are the square of the normalization constant divided by 2 (for wave functions initially normalized with their maximum value equal to unity), and approaches unity if the approximation (B14) is valid.

The proof of the second relation (B21) follows from application of the bound state equation

$$\begin{aligned}
& \sum_n \left(\frac{m_{01}^2}{z} + \frac{m_{02}^2}{1-z} - M_n^2 \right) \Phi_n(z, r_n) \int_0^1 dz' \Phi_n(z', r_n) \\
&= \sum_n \int_0^1 dy \left[\frac{\Phi_n(y, r_n) - \Phi_n(z, r_n)}{(y-z)^2} \right] \int_0^1 dz' \Phi_n(z', r_n) \\
&= \sum_n \left\{ \int_0^{z-\epsilon} + \int_{z-\epsilon}^{z+\epsilon} + \int_{z+\epsilon}^1 \right\} dy \left[\frac{\Phi_n(y, r_n) - \Phi_n(z, r_n)}{(y-z)^2} \right] \int_0^1 dz' \Phi_n(z', r_n) \\
&\rightarrow \sum_n \frac{1}{2} \int_{z-\epsilon}^{z+\epsilon} dy \Phi_n''(z, r_n) \int_0^1 dz' \Phi_n(z', r_n) = \\
&= \epsilon \sum_n \Phi_n''(z, r_n) \int_0^1 dz' \Phi_n(z', r_n) \rightarrow 0,
\end{aligned} \tag{B22}$$

where the identity (B20) is used to eliminate the nonsingular integrals over the intervals $[0, z - \epsilon]$ and $[z + \epsilon, 1]$, and the singular part of the integral is proportional to ϵ which vanishes as $\epsilon \rightarrow 0$.

If Eq. (2.41) is used, modifications of the completeness relation (2.51) and the identity (B20), which hold in the region $z \notin [0, 1]$, can be derived. The results are

$$\sum_n \left(\frac{m_1^2}{z} + \frac{m_2^2}{1-z} - M_n^2 \right) \Phi_n(z, r_n) \Phi_n(z', r_n) = \frac{g^2/\pi}{(z-z')^2} \quad z' \in [0, 1] \quad (\text{B23})$$

$$\sum_n \left(\frac{m_1^2}{z} + \frac{m_2^2}{1-z} - M_n^2 \right) \Phi_n(z, r_n) \int_0^1 dz' \Phi_n(z', r_n) = -\frac{g^2/\pi}{z(1-z)}. \quad (\text{B24})$$

Note that the completeness relation and all of these identities hold for *any* two quark Hamiltonian. In particular, the Hamiltonian may describe bound states with flavor (where, in general, $m_1 \neq m_2$) as well as flavorless bound states where $m_1 = m_2$. In some applications we will study the coupling of a photon to a $q\bar{q}$ pair. Such a coupling necessarily involves *flavorless* states only, and we will use the identities for the flavorless states.

APPENDIX C: SUMMATION OF THE QUARK SCATTERING SERIES

Here we show how to sum this series (2.58) and obtain a useful form for M .

Since V is independent of the plus components of momentum, the integrations over the plus components of momentum can be carried out. Using the identity (A7) [with $q_- < 0$ in this case], the first term on the rhs of (2.58) becomes

$$\begin{aligned} \langle MG\mathcal{O} \rangle_1(p'; q) &= i \int \frac{d^2 p}{\pi^2} \frac{V(p', p) \mathcal{O}(p_-, q)}{d_2(p_-, p_+, q)} \\ &= \int_0^1 dz V(z', z) \frac{\mathcal{O}(p_-, q)}{\Delta(z, q^2)} - \frac{g^2}{\pi z'(1-z')} \frac{\mathcal{O}(p'_-, q)}{\Delta(z', q^2)} \\ &= \int_0^1 dz \left\{ V(z', z) - \frac{g^2 \delta(z' - z)}{\pi z(1-z)} \right\} \frac{\mathcal{O}(p_-, q)}{\Delta(z, q^2)} \quad \text{if } z' \in [0, 1], \end{aligned} \quad (\text{C1})$$

or

$$\langle MG\mathcal{O} \rangle_1(p'; q) = i \int \frac{d^2 p}{\pi^2} \frac{V(p', p) \mathcal{O}(p_-, q)}{d_2(p_-, p_+, q)} = \int_0^1 dz V_0(z', z) \frac{\mathcal{O}(p_-, q)}{\Delta(z, q^2)} \quad \text{if } z' \notin [0, 1]. \quad (\text{C2})$$

Here the replacement (A9) has been used to convert $V \rightarrow V$ (or V_0). Similarly, the second term becomes

$$\begin{aligned} \langle MG\mathcal{O} \rangle_2(p'; q) &= i \int \frac{d^2 k}{\pi^2} \frac{V(p', k)}{d_2(k_-, k_+, q)} \int_0^1 dz \left\{ V(y, z) - \frac{g^2 \delta(y - z)}{\pi z(1-z)} \right\} \frac{\mathcal{O}(p_-, q)}{\Delta(z, q^2)} \\ &= \int_0^1 dy \left\{ V(z', y) - \frac{g^2 \delta(z' - y)}{\pi y(1-y)} \right\} \frac{1}{\Delta(y, q^2)} \\ &\quad \times \int_0^1 dz \left\{ V(y, z) - \frac{g^2 \delta(y - z)}{\pi z(1-z)} \right\} \frac{\mathcal{O}(p_-, q)}{\Delta(z, q^2)} \quad \text{if } z' \in [0, 1], \end{aligned} \quad (\text{C3})$$

or

$$\begin{aligned} \langle MGO \rangle_2(p'; q) &= \int_0^1 dy V_0(z', y) \frac{1}{\Delta(y, q^2)} \\ &\times \int_0^1 dz \left\{ V(y, z) - \frac{g^2 \delta(y - z)}{\pi z(1 - z)} \right\} \frac{\mathcal{O}(p_-, q)}{\Delta(z, q^2)} \quad \text{if } z' \notin [0, 1], \end{aligned} \quad (C4)$$

and the third term follows the same pattern. Replacing $(q_-)^2 M(p', p; q)/\pi \rightarrow M(z', z; q^2)$ and $\mathcal{O}(p_-, q) \rightarrow \mathcal{O}(z, q^2)$, the series (2.58), in the interval $z' \in [0, 1]$, is clearly summed by the following equation

$$\begin{aligned} \int_0^1 dz M(z', z; q^2) \frac{\mathcal{O}(z, q^2)}{\Delta(z, q^2)} &= \int_0^1 dz \left\{ V(z', z) - \frac{g^2 \delta(z' - z)}{\pi z(1 - z)} \right\} \frac{\mathcal{O}(z, q^2)}{\Delta(z, q^2)} \\ &+ \int_0^1 dy \frac{M(z', y; q^2)}{\Delta(y, q^2)} \int_0^1 dz \left\{ V(y, z) - \frac{g^2 \delta(y - z)}{\pi z(1 - z)} \right\} \frac{\mathcal{O}(z, q^2)}{\Delta(z, q^2)} \quad \text{if } z' \in [0, 1]. \end{aligned} \quad (C5)$$

Introducing the shorthand notation

$$F(z', z; q^2) = \frac{M(z', z; q^2)}{\Delta(z, q^2)}, \quad (C6)$$

Eq. (C5) can be rearranged into the following form

$$\begin{aligned} \int_0^1 dz F(z', z; q^2) &\left\{ \left(\Delta(z, q^2) + \frac{g^2/\pi}{z(1 - z)} \right) \frac{\mathcal{O}(z, q^2)}{\Delta(z, q^2)} - \int_0^1 dy V(z, y) \frac{\mathcal{O}(y, q^2)}{\Delta(y, q^2)} \right\} \\ &= \int_0^1 dz F(z', z; q^2) (H(z) - q^2) \frac{\mathcal{O}(z, q^2)}{\Delta(z, q^2)} \\ &= \int_0^1 dz \left\{ V(z', z) - \frac{g^2 \delta(z' - z)}{\pi z(1 - z)} \right\} \frac{\mathcal{O}(z, q^2)}{\Delta(z, q^2)}, \end{aligned} \quad (C7)$$

which requires that F satisfy the equation

$$F(z', z, q^2) (H(z) - q^2) = V(z', z) - \frac{g^2 \delta(z' - z)}{\pi z(1 - z)}. \quad (C8)$$

This equation is easily solved with the use of the Greens function:

$$\begin{aligned} F(z', z; q^2) &= - \int_0^1 dy \left\{ V(z', y) - \frac{g^2 \delta(z' - y)}{\pi y(1 - y)} \right\} G(y, z, q^2) \\ &= - \int_0^1 dy \left\{ V(z', y) - \frac{g^2 \delta(z' - y)}{\pi y(1 - y)} \right\} \sum_n \frac{\Phi_n(y, r_n) \Phi_n(z, r_n)}{q^2 - M_n^2} \\ &= \sum_n \left(M_n^2 - \frac{m_1^2}{z'} - \frac{m_2^2}{1 - z'} \right) \frac{\Phi_n(z', r_n) \Phi_n(z, r_n)}{q^2 - M_n^2}. \end{aligned} \quad (C9)$$

Putting it all together gives Eq. (2.59).

If $z' \notin [0, 1]$, the series reduces to the same formal result

$$\begin{aligned}
\int_0^1 dz M(z', z; q^2) \frac{O(z, q^2)}{\Delta(z, q^2)} &= \int_0^1 dz V_0(z', z) \frac{O(z, q^2)}{\Delta(z, q^2)} \\
&+ \int_0^1 dy V_0(z', y) \frac{1}{\Delta(y, q^2)} \int_0^1 dz M(y, z; q^2) \frac{O(z, q^2)}{\Delta(z, q^2)} \\
&= \int_0^1 dz V_0(z', z) \frac{O(z, q^2)}{\Delta(z, q^2)} + \sum_n \int_0^1 dy V_0(z', y) \\
&\quad \times \frac{\Delta(y, M_n^2)}{\Delta(y, q^2)} \frac{\Phi_n(y, r_n)}{M_n^2 - q^2} \int_0^1 dz \Phi_n(z, r_n) O(z, q^2) \\
&= \sum_n \Delta(z', M_n^2) \frac{\Phi_n(z', r_n)}{M_n^2 - q^2} \int_0^1 dz \Phi_n(z, r_n) O(z, q^2), \quad (C10)
\end{aligned}$$

where the completeness relation (2.51) and Eq. (2.41) were used in the last step. This shows that (2.59) can be applied for *any* $z' \in [-\infty, \infty]$.

APPENDIX D: SUMMATION OF THE QUARK CURRENT SERIES

The series (3.7) for the dressed quark current is evaluated in this Appendix. The “first” integral over k_+ in each term in the series is evaluated as in Eq. (C1) or (C2)

$$\begin{aligned}
i \int \frac{d^2 k}{(2\pi)^2} V(p', k) \gamma_- S_i(k) \gamma^\mu S_i(k - q) \gamma_- &\equiv i \int \frac{d^2 k}{\pi^2} \frac{V(p', k) N_i^\mu(k_-, q_-)}{d_2^i(k_-, k_+, q)} \\
&= \begin{cases} \int_0^1 dy \left\{ V(z', y) - \frac{g^2 \delta(z' - y)}{\pi y(1 - y)} \right\} \frac{N_i^\mu(k_-, q_-)}{\Delta_i(y, q^2)} & \text{if } z' \in [0, 1] \\ \int_0^1 dy V_0(z', y) \frac{N_i^\mu(k_-, q_-)}{\Delta_i(y, q^2)} & \text{if } z' \notin [0, 1], \end{cases} \quad (D1)
\end{aligned}$$

where

$$N_i^\mu(k_-, q_-) = \frac{\gamma_- [m_{0i} + k_- \gamma_+] \gamma^\mu [m_{0i} + (k_- - q_-) \gamma_+] \gamma_-}{4 k_- (k_- - q_-)}, \quad (D2)$$

and d_2^i and Δ_i are (A3) and (A6) with equal masses $m_1 = m_2 = m_i$. Note that the only dependence of the integrand on k_+ was in the denominator d_2^i , so the k_+ integral could be evaluated using the methods of Appendix C. The second integral is similarly evaluated. If $z' \in [0, 1]$, the following result is obtained for the series Eq. (3.7)

$$\begin{aligned}
j_i^\mu(p', p' - q) &= j_i^\mu(z', q^2) = \gamma^\mu + \int_0^1 dy \left\{ V(z', y) - \frac{g^2 \delta(z' - y)}{\pi y(1 - y)} \right\} \frac{N_i^\mu(y, q)}{\Delta_i(y', q^2)} \\
&+ \int_0^1 dy' \left\{ V(z', y') - \frac{g^2 \delta(z' - y')}{\pi y'(1 - y')} \right\} \frac{1}{\Delta_i(y', q^2)} \int_0^1 dy \left\{ V(y', y) - \frac{g^2 \delta(y' - y)}{\pi y(1 - y)} \right\} \frac{N_i^\mu(y, q)}{\Delta_i(y, q^2)} \\
&+ \dots = \gamma^\mu + \int_0^1 dz M(z', y; q^2) \frac{N_i^\mu(y, q)}{\Delta_i(y, q^2)}, \quad (D3)
\end{aligned}$$

where $N_i^\mu(y, q) = N_i^\mu(k_-, q_-)$, and the series is summed using Eq. (C5) for the scattering matrix M . Using the solution (2.59) for the scattering matrix the result, for all values of $z' = p'_-/q_-$, is

$$j_i^\mu(z', q^2) = \gamma^\mu + \sum_n \Delta_i(z', M_n^2) \frac{\Phi_n(z', r_n)}{M_n^2 - q^2} \int_0^1 dz \Phi_n(z, r_n) N_i^\mu(z, q), \quad (D4)$$

where $j_i^\mu(p', p' - q) = j_i^\mu(z', q^2)$, and Δ_i is the Δ of Eq. (A6) with equal masses $m_1 = m_2 = m_i$.

The same expression also holds for $z' \notin [0, 1]$, but the derivation differs. As before, the full result can be deduced from the form of the first two terms of the series (3.7). Guided by (D1), and using the steps already displayed in Eq. (C10), we obtain

$$\begin{aligned} j_i^\mu(z', q^2) &= \gamma^\mu + \int_0^1 dy V_0(z', y) \frac{N_i^\mu(y, q)}{\Delta_i(y', q^2)} \\ &+ \int_0^1 dy' V_0(z', y') \frac{1}{\Delta_i(y', q^2)} \int_0^1 dy \left\{ V(y', y) - \frac{g^2 \delta(y' - y)}{\pi y(1 - y)} \right\} \frac{N_i^\mu(y, q)}{\Delta_i(y, q^2)} \\ &+ \dots = \gamma^\mu + \int_0^1 dz M(z', y; q^2) \frac{N_i^\mu(y, q)}{\Delta_i(y, q^2)} \\ &= \gamma^\mu + \sum_n \Delta_i(z', M_n^2) \frac{\Phi_n(z', r_n)}{M_n^2 - q^2} \int_0^1 dz \Phi_n(z, r_n) N_i^\mu(z, q). \end{aligned} \quad (D5)$$

We see, as we did with Eq. (2.59), that (D4) holds for *all* z' .

Equation (D4) expresses the dressed current as a sum over contributions from the bound states that couple to the photon, showing that vector dominance is a rigorous consequence of this model.

The structure of the numerator N_i^μ depends on the bare current:

$$\begin{aligned} \gamma_- : \quad & N_i^-(y, q) = \gamma_- \\ \gamma_+ : \quad & N_i^+(y, q) = -\gamma_- \frac{m_{0i}^2}{2q_-^2 y(1 - y)} \\ \gamma_x : \quad & N_i^x(y, q) = -\gamma_x \gamma_- \frac{m_{0i}}{2q_- y(1 - y)}. \end{aligned} \quad (D6)$$

Note the interesting structure of the correction term to the bare current γ_x , which will be of central importance in our discussion below. Substituting these results into (D4) gives the following

$$\begin{aligned} j_{i-}(z', q^2) &= \gamma_- \left\{ 1 + \sum_n \Delta_i(z', M_n^2) \frac{\Phi_n(z', r_n)}{M_n^2 - q^2} \int_0^1 dy \Phi_n(y, r_n) \right\} \\ j_{i+}(z', q^2) &= \gamma_+ - \gamma_- \frac{m_{0i}^2}{2q_-^2} \left\{ \sum_n \Delta_i(z', M_n^2) \frac{\Phi_n(z', r_n)}{M_n^2 - q^2} \int_0^1 dy \frac{\Phi_n(y, r_n)}{y(1 - y)} \right\} \\ j_x(z', q^2) &= \gamma_x - \gamma_x \gamma_- \frac{m_{0i}}{2q_-} \left\{ \sum_n \Delta_i(z', M_n^2) \frac{\Phi_n(z', r_n)}{M_n^2 - q^2} \int_0^1 dy \frac{\Phi_n(y, r_n)}{y(1 - y)} \right\}. \end{aligned} \quad (D7)$$

We organize these expressions by introducing the quark form factor, defined in Eq. (3.8). This immediately gives Eq. (3.9) for the j_- current. The other currents in (D7) can also

be written in terms of F_i . To this end note that the bound state equation for equal mass quarks implies that

$$M_n^2 \int_0^1 dy \Phi_n(y, r_n) = m_{0i}^2 \int_0^1 dy \frac{\Phi_n(y, r_n)}{y(1-y)}. \quad (\text{D8})$$

Hence the quark mass can be removed from the second term in j_+ , giving

$$\begin{aligned} g_i(z', q) &\equiv -\frac{m_{0i}^2}{2q_-^2} \left\{ \sum_n \Delta_i(z', M_n^2) \frac{\Phi_n(z', r_n)}{M_n^2 - q^2} \int_0^1 dy \frac{\Phi_n(y, r_n)}{y(1-y)} \right\} \\ &= \frac{q_+}{q_-} \left\{ \sum_n \frac{M_n^2}{Q^2} \Delta_i(z', M_n^2) \frac{\Phi_n(z', r_n)}{M_n^2 + Q^2} \int_0^1 dy \Phi_n(y, r_n) \right\} \\ &= -\frac{q_+}{q_-} F_i(z', Q^2) + \frac{q_+}{q_- Q^2} \left\{ \sum_n \Delta_i(z', M_n^2) \Phi_n(z', r_n) \int_0^1 dy \Phi_n(y, r_n) \right\}, \quad (\text{D9}) \end{aligned}$$

where $g_i(z', q)$ defined by this equation should not be confused with the two body Greens function or the bound state vertex function. If $z' \in [0, 1]$, the second term can be reduced using the identities (B20) and (B21)

$$\sum_n \left[\frac{m_{0i}^2}{z'(1-z')} - M_n^2 - \frac{g^2/\pi}{z'(1-z')} \right] \Phi_n(z', r_n) \int_0^1 dy \Phi_n(y, r_n) = -\frac{g^2}{\pi} \frac{1}{z'(1-z')}. \quad (\text{D10})$$

The same result is obtained directly from identity (B24) if $z' \notin [0, 1]$. Hence, for all z' ,

$$g_i(z', q) = -\frac{q_+}{q_-} F_i(z', Q^2) + \frac{g^2}{2\pi q_-^2} \frac{1}{z'(1-z')} = -\frac{q_+}{q_-} F_i(z', Q^2) - \frac{g^2}{2\pi} \frac{1}{p'_- p_-}. \quad (\text{D11})$$

Finally, the j_x component of the current can be similarly reduced. The second term in j_x is proportional to

$$\begin{aligned} H_i(z', q) &\equiv -\frac{m_{0i}}{2q_-} \left\{ \sum_n \Delta_i(z', M_n^2) \frac{\Phi_n(z', r_n)}{M_n^2 - q^2} \int_0^1 dy \frac{\Phi_n(y, r_n)}{y(1-y)} \right\} \\ &= -\frac{1}{2q_- m_{0i}} \left\{ \sum_n M_n^2 \Delta_i(z', M_n^2) \frac{\Phi_n(z', r_n)}{M_n^2 + Q^2} \int_0^1 dy \Phi_n(y, r_n) \right\} \\ &= -\frac{q_+}{m_{0i}} F_i(z', Q^2) - \frac{1}{2q_- m_{0i}} \left\{ \sum_n \Delta_i(z', M_n^2) \Phi_n(z', r_n) \int_0^1 dy \Phi_n(y, r_n) \right\}. \quad (\text{D12}) \end{aligned}$$

The part of the second term in $\{\}$ is the same constant derived above, and combining the results gives Eq. (3.11).

APPENDIX E: KINEMATICS FOR DIS

Kinematics for DIS in the c.m. frame are given in this Appendix. When Q^2 is very large, the components of P and q in this frame are (see Ref. [14])

$$q_+ = Q \sqrt{\frac{1-x}{2x}} \left[1 - \frac{M_0^2 x(1-2x)}{2Q^2(1-x)} + \dots \right]$$

$$\begin{aligned}
q_- &= -Q\sqrt{\frac{x}{2(1-x)}} \left[1 + \frac{M_0^2 x(1-2x)}{2Q^2(1-x)} + \dots \right] \\
P_+ &= \frac{M_0^2}{Q}\sqrt{\frac{x(1-x)}{2}} \left[1 - \frac{M_0^2 x(1-2x^2)}{2Q^2(1-x)} + \dots \right] \\
P_- &= \frac{Q}{\sqrt{2x(1-x)}} \left[1 + \frac{M_0^2 x(1-2x^2)}{2Q^2(1-x)} + \dots \right].
\end{aligned} \tag{E1}$$

The energy and momentum of particles 1 and 2 in the final state are

$$\begin{aligned}
p_z &= \pm \frac{Q}{2}\sqrt{\frac{1-x}{x}} \left[1 + \frac{x(M_0^2 - 2m_1^2 - 2m_2^2)}{2Q^2(1-x)} + \dots \right] \\
p_{10} &= \sqrt{m_1^2 + p_z^2} = \frac{Q}{2}\sqrt{\frac{1-x}{x}} \left[1 + \frac{x(M_0^2 + 2m_1^2 - 2m_2^2)}{2Q^2(1-x)} + \dots \right] \\
p_{20} &= \sqrt{m_2^2 + p_z^2} = \frac{Q}{2}\sqrt{\frac{1-x}{x}} \left[1 + \frac{x(M_0^2 - 2m_1^2 + 2m_2^2)}{2Q^2(1-x)} + \dots \right].
\end{aligned} \tag{E2}$$

There are two possibilities corresponding to the two terms in the current (4.10). The first term in the current will be large only if the momentum of particle 1 is in the direction of q_3 (i.e. $p_{1z} = |p_z| > 0$). [Choosing it in the opposite direction requires a very large momentum flow through the wave function, and is suppressed.] In this case

$$\begin{aligned}
p_{1+}^{(1)} &= Q\sqrt{\frac{1-x}{2x}} \left[1 + \frac{x(M_0^2 - 2m_2^2)}{2Q^2(1-x)} + \dots \right] \\
p_{1-}^{(1)} &= \sqrt{\frac{x}{2(1-x)}} \frac{m_1^2}{Q} \left[1 - \frac{x(M_0^2 - 2m_2^2)}{2Q^2(1-x)} + \dots \right] \\
p_{2+}^{(1)} &= \sqrt{\frac{x}{2(1-x)}} \frac{m_2^2}{Q} \left[1 - \frac{x(M_0^2 - 2m_1^2)}{2Q^2(1-x)} + \dots \right] \\
p_{2-}^{(1)} &= Q\sqrt{\frac{1-x}{2x}} \left[1 + \frac{x(M_0^2 - 2m_1^2)}{2Q^2(1-x)} + \dots \right],
\end{aligned} \tag{E3}$$

where the superscript (1) is a reminder that these relations hold only for the *first* term in the current. The momentum relations for the *second* term in the current (4.10) follow by interchanging 1 and 2 on both sides of these equations. We see that the first term in the current gives $p_{1+} \rightarrow Q$ and $p_{2+} \rightarrow 1/Q$, while the second term gives $p_{1+} \rightarrow 1/Q$ and $p_{2+} \rightarrow Q$. The two terms describe kinematically distinct regions of phase space, and their interference can safely be neglected when the total cross section is computed. Using the definitions (4.7) and (4.9), z and z' can be easily related to the Bjorken variable x . We have

$$\begin{aligned}
z &= 1 - \frac{p_{2-}^{(1)}}{P_-} \simeq 1 - (1-x) = x \\
z' &= 1 - \frac{p_{1-}^{(2)}}{P_-} \simeq 1 - (1-x) = x.
\end{aligned} \tag{E4}$$

APPENDIX F: THE BOUND STATE TRANSITION CURRENTS

This Appendix includes details of the evaluation of the transition currents shown in Fig. 10. The e_1 term will be evaluated first. The e_2 term can then be obtained by a simple substitution, as discussed at the end of this Appendix.

The e_1 term, given in Eq. (3.13), is reduced by introducing the momentum fractions

$$\xi = \frac{(P+k)_-}{P_-} \quad \eta = \frac{(P_f+k)_-}{q_-} \quad y = -\frac{q_-}{P_-}, \quad (\text{F1})$$

so that

$$\frac{P_{f-}}{P_-} = 1 - y \quad \frac{(P+k)_-}{q_-} = \eta - 1 \quad \xi' = \frac{(P_f+k)_-}{P_{f-}} = \frac{\xi - y}{1 - y}. \quad (\text{F2})$$

In this notation the x -type transverse quark current, evaluated in Eq. (3.11), is

$$\begin{aligned} j_1^x &= \gamma_x - \gamma_x \gamma_- \left\{ \frac{g^2}{2\pi m_{01}} \frac{q_-}{(P_{f-} + k_-)(P_- + k_-)} - \frac{q_+}{m_{01}} F_1(\eta, Q^2) \right\} \\ &= \gamma_x + \gamma_x \gamma_- H_1(\eta, q), \end{aligned} \quad (\text{F3})$$

First evaluate the numerators, N^μ , of the traces in (3.13). The numerator N_x is

$$\begin{aligned} N_x &= \text{Tr} \left[\gamma_- \gamma_x \left\{ m_{01} + (P_f + k)_- \gamma_+ \right\} (\gamma_x + \gamma_x \gamma_- H_1(\eta, q)) \right. \\ &\quad \left. \times \left\{ m_{01} + (P + k)_- \gamma_+ \right\} \gamma_- (k_- \gamma_+) \right] \\ &= 16 k_- [m_{01} q_- + 2(P_f + k)_- (P + k)_- H_1(\eta, q)] \\ &= 16 k_- q_- \frac{m_1^2}{m_{01}} - 32(P_f + k)_- (P + k)_- \frac{q_+}{m_{01}} F_1(\eta, Q^2) \\ &= \frac{16 k_- q_-}{m_{01}} \left(m_1^2 - \eta(1 - \eta) Q^2 F_1(\eta, Q^2) \right), \end{aligned} \quad (\text{F4})$$

where the trace has been evaluated in a Dirac space of four dimensions. The numerator of the minus component of the current has only one term

$$\begin{aligned} N_- &= \text{Tr} \left[\gamma_- (P_f + k)_- \gamma_+ \gamma_- (P + k)_- \gamma_+ \gamma_- (k_- \gamma_+) \right] [1 + F_1(\eta, Q^2)] \\ &= 32 k_- (P_f + k)_- (P + k)_- [1 + F_1(\eta, Q^2)] \\ &= 16 k_- \frac{q_-}{q_+} \eta(1 - \eta) Q^2 [1 + F_1(\eta, Q^2)]. \end{aligned} \quad (\text{F5})$$

The trace for the plus component is

$$\begin{aligned} N_+ &= \text{Tr} \left[\gamma_- \left\{ m_{01} + (P_f + k)_- \gamma_+ \right\} (\gamma_+ + \gamma_- G_1(\eta, q)) \right. \\ &\quad \left. \times \left\{ m_{01} + (P + k)_- \gamma_+ \right\} \gamma_- (k_- \gamma_+) \right] \\ &= 16 k_- m_1^2 - 32 k_- (P_f + k)_- (P + k)_- \frac{q_+}{q_-} F_1(\eta, Q^2) \\ &= 16 k_- \left(m_1^2 - \eta(1 - \eta) Q^2 F_1(\eta, Q^2) \right). \end{aligned} \quad (\text{F6})$$

Inserting the numerator (F5) into the general result (3.13) gives the following result for the minus component of the current

$$\begin{aligned}
& \langle f_- | \mathcal{J}_-(P_f, P) | i_- \rangle \Big|_{e_1 \text{ term}} \\
&= 32ie_1 \int \frac{d^2k}{(2\pi)^2} \frac{(P_f + k)_-(P + k)_- k_- G_f(-P_f - k, -k) G_i(P + k, k)}{[m_1^2 - (P_f + k)^2 - i\epsilon][m_1^2 - (P + k)^2 - i\epsilon][m_2^2 - k^2 - i\epsilon]} [1 + F_1(\eta, Q^2)] \\
&= 32ie_1 \int \frac{d^2k}{(2\pi)^2} \frac{1}{d_3} G_f(-P_f - k, -k) G_i(P + k, k) [1 + F_1(\eta, Q^2)] , \tag{F7}
\end{aligned}$$

where the subscript - on f_- reminds us that the final state must have a pure γ_- structure (if the ground state does), d_3 is the product of the denominators of the three quark propagators, with three poles in k_+

$$d_3 = \left[\frac{m_1^2}{\xi' P_{f-}} - \frac{M_f^2}{P_{f-}} - 2k_+ - i\epsilon_a \right] \left[\frac{m_1^2}{\xi P_-} - \frac{M_i^2}{P_-} - 2k_+ - i\epsilon_b \right] \left[\frac{m_2^2}{(\xi - 1) P_-} - 2k_+ - i\epsilon_c \right] , \tag{F8}$$

and the ϵ 's change sign according to

$$\begin{aligned}
\epsilon_a &= \epsilon / (P_f + k)_- = \epsilon / (\xi' P_{f-}) = \epsilon / [(\xi - y) P_-] \\
\epsilon_b &= \epsilon / (P + k)_- = \epsilon / (\xi P_-) \\
\epsilon_c &= \epsilon / k_- = \epsilon / [(\xi - 1) P_-] . \tag{F9}
\end{aligned}$$

Since the vertex functions G do not depend on k_+ , we can evaluate the k_+ integral. It will be nonzero only when the three poles of (F8) do not all lie in the same half plane. Since $q_- < 0$ for electron scattering, $P_- > P_{f-} > 0$, which implies that

$$0 < y < 1 . \tag{F10}$$

Hence

$$\xi - 1 < \xi - y < \xi , \tag{F11}$$

and all three poles will be in the same half of the complex plane (giving zero for the integral) unless

$$0 < \xi < 1 . \tag{F12}$$

There are two terms, depending on the sign of $\xi - y$. Closing the k_+ contour in the upper half plane, and using $dk_- = P_- d\xi$ and $(\xi - 1)P_- = (\xi' - 1)P_{f-}$ gives

$$\begin{aligned}
\langle f_- | \mathcal{J}_-(P_f, P) | i_- \rangle \Big|_{e_1 \text{ term}} &= 16e_1 P_-^2 P_{f-} \int_0^1 \frac{d\xi}{(2\pi)} \frac{G_f(\xi', P_f) G_i(\xi, P)}{\Delta(\xi', P_f^2) \Delta(\xi, P^2)} [1 + F_1(\eta, Q^2)] \\
&\quad - 16e_1 P_-^2 P_{f-} \int_0^y \frac{d\xi}{(2\pi)} \frac{G_f(\xi', P_f) G_i(\xi, P) [1 + F_1(\eta, Q^2)]}{\Delta(\xi', P_f^2) \left(\Delta(\xi, P^2) - \frac{\Delta(\xi', P_f^2)}{(1 - y)} \right)} , \tag{F13}
\end{aligned}$$

where the first term is the contribution from the pole due to the zero in the third term in (F8) above, and the second from the pole due to the zero in the first term. Using the

definition (2.39) of the wave function and combining the two contributions in the region $[0, y]$ we get Eq. (3.14) with

$$\begin{aligned}\mathcal{R} &= \frac{\Delta(\xi', P_f^2)}{\Delta(\xi', P_f^2) - (1-y)\Delta(\xi, P^2)} \\ &= \frac{\xi \left[\xi'(1-\xi')M_f^2 - (1-\xi')m_1^2 - \xi'm_2^2 \right]}{(1-\xi') \left\{ \xi \xi' (M_f^2 - (1-y)M_i^2) - y m_1^2 \right\}} \rightarrow 1 \quad \text{if } M_f^2 \rightarrow \infty. \quad (\text{F14})\end{aligned}$$

This factor is needed for positive values of ξ in the interval $[0, y]$ and *negative* values of ξ' in the interval $[-y/(1-y), 0]$. In this region the denominator has a zero only if $m_1^2 < 0$. In order to avoid the discussion of such cases we limit numerical applications to cases with $m_1^2 > 0$.

Results for the other components of the transition current, Eq. (3.16), are obtained by a similar argument using (F4) and (F6) in place of (F5).

Now consider the modifications required in order to evaluate the e_2 term. Using the momenta defined in Fig. 10, and the same definitions of momentum fractions (F1) and (F2), the e_2 term is obtained from the e_1 term simply by substituting $m_1 \leftrightarrow m_2$ and $e_1 \rightarrow e_2$. However, the momentum fractions in the wave functions are, by convention, the fraction of the momentum carried by the quark m_1 , and the momentum fraction in the quark form factor is that of the *outgoing* quark, and hence these functions must be written in terms of

$$\begin{aligned}\xi_2 &= \frac{-k_-}{P_-} = 1 - \xi \\ \xi'_2 &= \frac{-k_-}{P_{f-}} = 1 - \xi' \\ \eta_2 &= \frac{-(P+k)_-}{q_-} = 1 - \eta, \quad (\text{F15})\end{aligned}$$

These observations lead immediately to the final result (3.17).

APPENDIX G: THE MODIFIED CUBIC SPLINES

The two body equations were solved using a modification of the standard cubic splines employed previously in many problems. The standard splines are defined on 4 segments of length h , bounded by the 5 points $a = (n-2)h, b = a + h, c = b + h, d = c + h, e = d + h$:

$$S_n(x) = \frac{1}{4} \begin{cases} \frac{(x-a)^3}{h^3} & \text{if } a < x < b \\ 1 + 3 \left[\frac{(x-b)}{h} + \frac{(x-b)^2}{h^2} - \frac{(x-b)^3}{h^3} \right] & \text{if } b < x < c \\ 1 + 3 \left[\frac{(d-x)}{h} + \frac{(d-x)^2}{h^2} - \frac{(d-x)^3}{h^3} \right] & \text{if } c < x < d \\ \frac{(e-x)^3}{h^3} & \text{if } d < x < e. \end{cases} \quad (\text{G1})$$

If the interval $[0, 1]$ is spanned by $n_s - 2$ standard splines, there must be $n_s + 1$ segments, of length $h = 1/(n_s + 1)$. The first spline is numbered $n = 2$ beginning at $x = 0$ and the last is numbered $n_s - 1$ ending at $x = 1$.




FIG. 14: The modified cubic splines for the minimal case of $n_s = 3$, with four segments. In the left figure the left- and right-most splines at the boundary approach 0 as $x^{0.6705}$ and $(1-x)^{0.6705}$ appropriate to quark masses $m_{01} = m_{02} = 1.5$. In the right figure the bare quark masses are zero and the splines approach a constant at the boundaries. Note that the sum of the splines (the dotted line) is a constant in the central region, and throughout the *whole* region in the chiral case.

The modified splines used in this paper consist of the standard splines plus one additional, non-standard spline inserted at the beginning and end of the interval $[0,1]$. These non-standard splines are defined on only three segments, and will be numbered 1 and n_s . They are

$$\begin{aligned}
 S_1(x) &= \begin{cases} d_1 \frac{x^{\beta_1}}{h^{\beta_1}} + d_2 \frac{x^2}{h^2} + d_3 \frac{x^3}{h^3} & \text{if } 0 < x < h \\ \frac{1}{4} + \frac{3}{4} \left[\frac{(2h-x)}{h} + \frac{(2h-x)^2}{h^2} - \frac{(2h-x)^3}{h^3} \right] & \text{if } h < x < 2h \\ \frac{(3h-x)^3}{4h^3} & \text{if } 2h < x < 3h \end{cases} \\
 S_{n_s}(x) &= \begin{cases} \frac{(x-1+3h)^3}{4h^3} & \text{if } 1-3h < x < 1-2h \\ \frac{1}{4} + \frac{3}{4} \left[\frac{(1-h-x)}{h} + \frac{(1-h-x)^2}{h^2} - \frac{(1-h-x)^3}{h^3} \right] & \text{if } 1-2h < x < 1-h \\ d_1 \frac{(1-x)^{\beta_2}}{h^{\beta_2}} + d_2 \frac{(1-x)^2}{h^2} + d_3 \frac{(1-x)^3}{h^3} & \text{if } 1-h < x < 1 \end{cases} \quad (\text{G2})
 \end{aligned}$$

where the β_i are the fractional exponents determined by the boundary conditions [given in Eq. (2.49)], and the coefficients d_i depend on the exponents β

$$d_1 = \frac{3}{(3-\beta)(2-\beta)}, \quad d_2 = \frac{3(1-\beta)}{(2-\beta)}, \quad d_3 = \frac{(2\beta-3)}{(3-\beta)}, \quad (\text{G3})$$

where $\beta = \beta_1$ for S_1 or β_2 for S_{n_s} . These coefficients were fixed by the requirement that the spline and its first two derivatives be continuous. Figure 14 shows two examples of the modified splines for the minimal number $n_s = 3$. In this case the central spline spans the full interval $[0,1]$ and has 4 segments.

Using these splines as a basis, the equation is reduced to a matrix equation that is solved with the standard eigenvalue subroutine packages.

-
- [1] G. 't Hooft, Nucl. Phys. **B75**, 461 (1974).
 - [2] C. G. Callan, N. Coote and D. J. Gross, Phys. Rev. D **13**, 1649 (1976).
 - [3] M. Einhorn, Phys. Rev. D **14**, 3451 (1976).
 - [4] M. I Bars, M. B. Green, Phys. Rev. D **15**, (1978).
 - [5] C. G. Callan and D. J. Gross, Phys. Rev. Lett. **22**, 156 (1969).
 - [6] M. Burkardt, Phys. Rev. D **62**, 094003 (2000) [arXiv:hep-ph/0005209].
 - [7] M. Burkardt and N. Uraltsev, Phys. Rev. D **63**, 014004 (2001) [arXiv:hep-ph/0005278].
 - [8] V. Schon and M. Thies, Contribution to the Festschrift in honor of Boris Ioffe, M. Shifman, (ed). In *At the frontier of particle physics, vol. 3* p. 1945 (2000) [arXiv:hep-th/0008175].
 - [9] M. Burkardt, F. Lenz and M. Thies, Phys. Rev. D **65**, 125002 (2002) [arXiv:hep-th/0201235].
 - [10] Y. S. Kalashnikova and A. V. Nefediev, Phys. Usp. **45**, 347 (2002) [Usp. Fiz. Nauk **172**, 378 (2002)] [arXiv:hep-ph/0111225].
 - [11] B. Grinstein, R. F. Lebed, Phys. Rev. D **57**:1366 (1998) [hep-ph/9708396].
 - [12] R. F. Lebed, Talk given at 3rd International Conference in Quark Confinement and Hadron Spectrum (Confinement III), Newport News, VA, 7-12 Jun 1998; Published in *Newport News 1998, Quark confinement and the hadron spectrum III*, p.161 (1998) [hep-ph/9808452].
 - [13] R. F. Lebed, N. G. Uraltsev, Phys. Rev. D **62**:094011 (2000) [hep-ph/0006346].
 - [14] Z. Batiz and F. Gross, Phys. Rev. C **58**, 2963 (1998).
Note that the square bracket in the second of Eqs. (5.8) should be $\left[1 - \frac{M^2 x(1-2x)^2}{2Q^2(1-x)}\right]$ instead of $\left[-\frac{M^2 x(1-2x)^2}{2Q^2(1-x)}\right]$.
 - [15] Z. Batiz and F. Gross, in preparation.
 - [16] F. Gross and J. Milana, Phys. Rev. D **43**(1991) 2401.
F. Gross and J. Milana, Phys. Rev. D **45** (1992) 969.
F. Gross and J. Milana, Phys. Rev. D **50** (1994) 3332.
 - [17] C. Savkli and F. Gross, Phys. Rev. C **63**, 035208 (2001).
 - [18] F. Gross and D. O. Riska, Phys. Rev. C **36**, 1928 (1987).
 - [19] F. E. Close, N. Isgur, Phys. Lett. B **509**, 81 (2001) [hep-ph/0102067].
 - [20] N. Isgur, S. Jeschonnek, W. Melnitchouk and J. W. Van Orden, Phys. Rev. D **64**:054005 (2001).